

NUWC-NPT Technical Report 12,298
17 August 2020

Dive Distribution and Group Size Parameters for Marine Species Occurring in the U.S. Navy's Northwest Training and Testing Study Area

Stephanie L. Watwood
Jocelyn R. Borcuk
Emily R. Robinson
NUWC Division Newport

Erin M. Oliveira
McLaughlin Research Corporation

Stephanie L. Sleeman
Naval Facilities Engineering Command Northwest



Naval Undersea Warfare Center Division Newport, Rhode Island

Approved for public release; distribution is unlimited.

PREFACE

An important element of the Navy's comprehensive environmental planning is the acoustic effects analysis executed with the Navy Acoustic Effects Model (NAEMO) software. NAEMO was developed to estimate the possible impacts of anthropogenic sound on marine animals, combining established acoustic propagation modeling with data regarding the distribution and abundance of marine species. This report recommends species-typical static depth distributions and group size information for all marine mammal and sea turtle species that occur in the Northwest Training and Testing (NWTT) Study Area that will be modeled using NAEMO.

Technical reviewers for the information within this report were Chip Johnson (Command Pacific Fleet), Laura Busch (United States Fleet Forces), Andrew DiMatteo (McLaughlin Research Corporation), Dawn Grebner (Naval Undersea Warfare Center Division, Keyport), Jennifer Paulk (Naval Air Systems Command), and Tara Moll (Naval Undersea Warfare Center Division, Newport).

Reviewed and Approved: 29 October 2018



Dennis Tierney
Head, Infrastructure Division



REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188	
<p>The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OPM control number.</p> <p>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 29-10 2018		2. REPORT TYPE Technical Report		3. DATES COVERED (From – To) September 2018 – November 2020	
4. TITLE AND SUBTITLE Dive Distribution and Group Size Parameters for Marine Species Occurring in the U.S. Navy's Northwest Training and Testing Study Area				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER N/A	
				5c. PROGRAM ELEMENT NUMBER N/A	
6. AUTHOR(S) Stephanie L. Watwood, Jocelyn R. Borcuk, and Emily R. Robinson NUWC Division, Newport Erin M. Oliveira McLaughlin Research Corporation Stephanie Sleeman Naval Facilities Engineering Command Northwest				5.d PROJECT NUMBER N/A	
				5e. TASK NUMBER N/A	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Undersea Warfare Center Division 1176 Howell Street Newport, RI 02841-1708				8. PERFORMING ORGANIZATION REPORT NUMBER TR 12,298	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSORING/MONITOR'S ACRONYM	
				11. SPONSORING/MONITORING REPORT NUMBER	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT. An important element of the Navy's comprehensive environmental planning is the acoustic effects analysis executed with the Navy Acoustic Effects Model (NAEMO) software. NAEMO was developed to estimate the possible impacts of anthropogenic sound on marine animals, combining established acoustic propagation modeling with data regarding the distribution and abundance of marine species. This report recommends species-typical static depth distributions and group size information for all marine mammal and sea turtle species that occur in the Northwest Training and Testing (NWTT) Study Area that will be modeled using NAEMO.					
15. SUBJECT TERMS Navy Acoustic Effects Model (NAEMO), Acoustic modeling, Marine Mammals, Sea Turtles, Northwest Training and Testing (NWTT)					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Stephanie L. Watwood
U	U	U	SAR	77	19b. TELEPHONE NUMBER (Include area code) (401) 832-4929

This page left intentionally blank.

EXECUTIVE SUMMARY

An important element of the Navy's comprehensive environmental planning is the acoustic effects analysis executed with the Navy Acoustic Effects Model (NAEMO) software. NAEMO was developed to estimate the possible impacts of anthropogenic sound on marine animals, combining established acoustic propagation modeling with data regarding the distribution and abundance of marine species. This report recommends species-typical static depth distributions and group size information for all marine mammal and sea turtle species that occur in the Northwest Training and Testing (NWTT) Study Area that will be modeled using NAEMO.

This page left intentionally blank.

TABLE OF CONTENTS

Section	Page
EXECUTIVE SUMMARY	ES-1
LIST OF ABBREVIATIONS AND ACRONYMS	ii
1 INTRODUCTION	1-1
1.1 The Navy Acoustic Effects Model	1-1
1.2 Data Inputs	1-1
2 MARINE MAMMAL AND SEA TURTLE DEPTH DISTRIBUTIONS	2-1
2.1 Surrogate Species and Study Areas	2-1
2.2 Marine Mammal and Sea Turtle Dive Behavior Summaries	2-5
2.2.1 Cetaceans	2-5
2.2.2 Carnivores	2-25
2.2.3 Sea Turtles	2-31
2.3 Conclusions	2-32
3 MARINE MAMMAL GROUP SIZE INFORMATION	3-1
4 REFERENCES	4-1

LIST OF ILLUSTRATIONS

Figure	Page
1 Northwest Training and Testing Study Area	2-2

LIST OF TABLES

Table	Page
1 Marine Mammal and Sea Turtle Species Occurring in the NWTT Study Area	2-3
2 Percentage of Time at Depth for the Minke Whale ¹	2-6
3 Percentage of Time at Depth for the Bryde's Whale ^{1,2}	2-7
4 Percentage of Time at Depth for the Blue Whale ¹	2-8
5 Percentage of Time at Depth for the Fin Whale ¹	2-9
6 Percentage of Time at Depth for the Humpback Whale on Foraging Grounds ¹	2-10
7 Percentage of Time at Depth for the Short-Finned Pilot Whale ^{1,2}	2-11

LIST OF TABLES (Cont'd)

Table	Page
8	Percentage of Time at Depth for the Risso's Dolphin ¹ 2-12
9	Percentage of Time at Depth for Fish-Eating Killer Whales ¹ 2-13
10	Percentage of Time at Depth for Mammal-Eating Killer Whales ¹ 2-13
11	Percentage of Time at Depth for the Pantropical Spotted Dolphin ^{1,2} 2-15
12	Percentage of Time at Depth for the Bottlenose Dolphin ¹ 2-16
13	Percentage of Time at Depth for the Gray Whale ¹ 2-17
14	Percentage of Time at Depth for the Harbor Porpoise ¹ 2-19
15	Percentage of Time at Depth for the Dall's Porpoise ¹ 2-20
16	Percentage of Time at Depth for the Sperm Whale in the Pacific Ocean ¹ 2-21
17	Percentage of Time at Depth for the Blainville's Beaked Whale ^{1,2} 2-23
18	Percentage of Time at Depth for the Cuvier's Beaked Whale ^{1,2} 2-24
19	Percentage of Time at Depth for the Northern Sea Otter 2-26
20	Percentage of Time at Depth for the Guadalupe Fur Seal ¹ 2-27
21	Percentage of Time at Depth for the Northern Fur Seal ¹ 2-27
22	Dive Depth Distribution Data from Rehberg et al. (2009), Merrick (1995), and Merrick and Loughlin (1997) Used to Estimate Depth Distribution for the Steller Sea Lion 2-29
23	Percentage of Time at Depth for the Stellar Sea Lion ¹ 2-29
24	Percentage of Time at Depth for the California Sea Lion ¹ 2-30
25	Percentage of Time at Depth for the Northern Elephant Seal ¹ 2-31
26	Percentage of Time at Depth for the Harbor Seal ¹ 2-32
27	Percentage of Time at Depth for the Leatherback Turtle ¹ 2-34
28	Mean Group Size, Standard Deviation, and Ranges for Marine Species in the Offshore Waters Portion of the NWTT Study Area 3-1
29	Mean Group Size, Standard Deviation, and Range for Marine Species in the Inland Waters of Puget Sound and San Juan Islands/Strait of Juan de Fuca Portion of the NWTT Study Area 3-4
30	Mean Group Size, Standard Deviation, and Range for Marine Species in the Behm Canal Portion of the NWTT Study Area 3-5

LIST OF ABBREVIATIONS AND ACRONYMS

DTAG	Digital Acoustic Recording Tag
E AK	Eastern Alaska
M	Meter(s)
Min	Minute(s)
°N	Degrees North
NAEMO	Navy Acoustic Effects Model
NMFS	National Marine Fisheries Services
NUWC	Naval Undersea Warfare Center
NWTT	Northwest Training and Testing
SD	Standard Deviation
TDR	Time-Depth Recorder
U.S.	United States
USFWS	U.S. Fish and Wildlife Services
W AK	Western Alaska

This page left intentionally blank.

1. INTRODUCTION

The United States (U.S.) Navy is required to assess potential impacts of Navy-generated sound in the water on protected marine species in compliance with applicable laws and regulations, including the National Environmental Policy Act, Executive Order 12114, the Marine Mammal Protection Act, and the Endangered Species Act. This report describes the methods and analytical approach to quantifying the depth distributions in the water column and group sizes of marine mammals and sea turtles to be used within the Navy Acoustic Effects Model (NAEMO).

1.1 The Navy Acoustic Effects Model

NAEMO is the standard model used by the Navy to estimate potential impacts to marine species from impulsive and non-impulsive sound sources used during Navy training and testing activities. NAEMO combines marine species distribution information with environmental parameters, propagation characteristics, sound source parameters, and typical training or testing scenarios in order to assess the level of behavioral disturbance, hearing impacts (including both temporary and permanent threshold shifts), and other injuries predicted for individual marine mammals and sea turtles likely to be in the vicinity of Navy training and testing activities.

1.2 Data Inputs

NAEMO first uses location-specific density (more detailed information regarding species density is available in density technical reports by U.S. Department of the Navy (DiMatteo et al. 2017; Hanser et al. 2017)) and group size information to patchily distribute a given marine species into a simulation area. The depth distribution data are then used to place animals in the water column at the depths at which they are typically found. An animal is reassigned a new depth every four minutes throughout the simulation, based on the depth distribution for that species. When available, seasonal or geographically-specific depth and group size information is used.

In addition to available marine mammal and sea turtle data, specific information about environmental conditions and projected Navy activities within a study area is needed to run NAEMO and quantify potential impacts to marine mammals and sea turtles. This environmental data include information about bathymetry, seafloor composition (e.g., rock, sand), and factors that vary throughout the year such as wind speed and sound velocity profiles. The details of Navy training and testing activities are also collected, including location, frequency, and source characteristics. For more detailed information about the NAEMO model, consult the Quantitative Analysis Technical Report (U.S. Department of the Navy 2018).

This page left intentionally blank.

2. Marine Mammal and Sea Turtle Depth Distributions

The best available science from literature reviews was used to obtain species-specific depth distribution information for the Northwest Training and Testing Study Area (NWTT; Figure 1). Journal articles, books, technical reports, cruise reports, funding agency reports, theses, dissertations, and raw data from individual researchers were assessed for this report.

As described in Section 1.2, depth distribution data are combined with species density data during the NAEMO modeling process. Densities were obtained from the Navy Marine Species Density Database (DiMatteo et al. 2017; Hanser et al. 2017). In some instances, density data were provided for guilds of species, rather than for individual species (e.g., the small beaked whale guild is a group of multiple species). This multi-species guild was created because observers could not differentiate between closely related species at sea, or because the sample sizes of the species observed individually were too small to incorporate into density modeling. For this case, a single representative species was chosen to correspond to the multi-species groupings contained in the density data.

The information required for representing a species in NAEMO specifically focuses on the percent of time each animal spends in the water column, defined here as a range of depths extending from the surface to the maximum dive depth of each species. Depth distributions contain percent time spent in the water only, either at the surface or in given depth bins. Percentage values may be slightly above or below 100 percent, due to decimal rounding, especially when animals spend smaller percentages of time close to their maximum dive depth. Rather than round down to zero, the deep bins are often rounded up in order to show a fraction of a percentage when it has been recorded that a species is capable of reaching that depth bin. For pinniped species, time spent hauled out of the water is not represented (this is accounted for in the density data).

2.1 Surrogate Species and Study Areas

Depth distribution data within this report are based upon species-specific tagging data obtained during literature review. If tagging data were not available for a particular species, data for the most similar species were used in the form of a surrogate. A species will generally only be considered a surrogate for modeling if the species is closely related (i.e., within the same genus or family), feeds on similar prey, or has a distribution in similar water types (e.g., continental shelf waters). The exception to the general surrogate selection is the two species of *Kogia* spp. (dwarf and pygmy sperm whales), for which there are no other species in their family to choose as surrogates. Therefore, a species from another family within their suborder (Odontoceti) was chosen as a surrogate. Surrogate species (if required) for all species are provided in Table 1 for the NWTT Study Area.




Figure 1. Northwest Training and Testing Study Area

Table 1. Marine Mammal and Sea Turtle Species Occurring in the NWTT Study Area

Species Name	Common Name	Surrogate species	Offshore	Inland Waters	Behm Canal	Section
Cetaceans						
Family Balaenopteridae						
<i>Balaenoptera acutorostrata</i>	Common minke whale	N/A	X	X	X	2.2.1.1.1
<i>Balaenoptera borealis</i>	Sei whale	Bryde's whale (<i>Balaenoptera edeni</i>)	X			2.2.1.1.2
<i>Balaenoptera edeni</i>	Bryde's whale	N/A				2.2.1.1.3
<i>Balaenoptera musculus</i>	Blue whale	N/A	X			2.2.1.1.4
<i>Balaenoptera physalus</i>	Fin whale	N/A	X		X	2.2.1.1.5
<i>Megaptera novaeangliae</i>	Humpback whale	N/A	X	X	X	2.2.1.1.6
Family Delphinidae						
<i>Delphinus delphis</i>	Short-beaked common dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X			2.2.1.2.1
<i>Globicephala macrorhynchus</i>	Short-finned pilot whale	N/A	X			2.2.1.2.2
<i>Grampus griseus</i>	Risso's dolphin	N/A	X			2.2.1.2.3
<i>Orcinus orca</i>	Killer whale	N/A	X (East Pacific Offshore, West Coast Transients, Southern Residents, Northern Residents)	X (Southern Residents, Northern Residents, East West Coast Transients)	X (Alaska Residents, West Coast Transients)	2.2.1.2.4
<i>Lagenorhynchus obliquidens</i>	Pacific white-sided dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X	X	X	2.2.1.2.5
<i>Lissodelphis borealis</i>	Northern right whale dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X			2.2.1.2.6

Table 1. Marine Mammal and Sea Turtle Species Occurring in the NWTT Study Area (Cont'd)

Species Name	Common Name	Surrogate species	Offshore	Inland Waters	Behm Canal	Section
Cetaceans						
Family Delphinidae						
<i>Stenella attenuata</i>	Pantropical spotted dolphin	N/A				2.2.1.2.7
<i>Stenella coeruleoalba</i>	Striped dolphin	Pantropical spotted dolphin (<i>Stenella attenuata</i>)	X			2.2.1.2.8
<i>Tursiops truncatus</i>	Common bottlenose dolphin	N/A	X			2.2.1.2.9
Family Eschrichtiidae						
<i>Eschrichtius robustus</i>	Gray whale	N/A	X	X		2.2.1.3.1
Family Kogiidae						
<i>Kogia spp.guild</i>	Pygmy and dwarf sperm whale	Short-finned pilot whale (<i>Globicephala macrorhynchus</i>)	X			2.2.1.4.1
Family Phocoenidae						
<i>Phocoena phocoena</i>	Harbor porpoise	N/A	X	X	X	2.2.1.5.1
<i>Phocoenoides dalli</i>	Dall's porpoise	N/A	X	X	X	2.2.1.5.2
Family Physeteridae						
<i>Physeter macrocephalus</i>	Sperm whale	N/A	X			2.2.1.6.1
Family Ziphiidae						
<i>Berardius bairdii</i>	Baird's beaked whale	Cuvier's beaked whale (<i>Ziphius cavirostris</i>)	X			2.2.1.7.1
<i>Mesoplodon densirostris</i>	Blainville's beaked whale	N/A				2.2.1.7.2
<i>Ziphius cavirostris</i>	Cuvier's beaked whale	N/A				2.2.1.7.3
<i>Small beaked whale guild</i>	<i>Ziphius</i> and <i>Mesoplodon</i> spp.	N/A	X			2.2.1.7.4

Table 1. Marine Mammal and Sea Turtle Species Occurring in the NWTT Study Area (Cont'd)

Species Name	Common Name	Surrogate species	Offshore	Inland Waters	Behm Canal	Section
Sea Otters						
Family Mustelidae						
<i>Enhydra lutris</i>	Northern Sea Otter	N/A	X			2.2.2.1.1
Pinnipeds						
Family Otariidae						
<i>Arctocephalus townsendi</i>	Guadalupe fur seal	N/A	X			2.2.2.2.1
<i>Callorhinus ursinus</i>	Northern fur seal	N/A	X		X	2.2.2.2.2
<i>Eumetopias jubatus</i>	Steller sea lion	N/A	X	X	X	2.2.2.2.3
<i>Zalophus californianus</i>	California sea lion	N/A	X	X		2.2.2.2.4
Family Phocidae						
<i>Mirounga angustirostris</i>	Northern elephant seal	N/A	X	X	X	2.2.2.3.1
<i>Phoca vitulina</i>	Harbor seal	N/A	X	X	X	2.2.2.3.2
Sea Turtles						
Family Dermochelyidae						
<i>Dermochelys coriacea</i>	Leatherback sea turtle	N/A	X			2.2.3.1.1

2.2 Marine Mammal and Sea Turtle Dive Behavior Summaries

This section discusses the depth distributions that were constructed for each species or surrogate species, based on the best available science. Ideally, depth distributions would be specific to different locations; however, sometimes diving data were not available for the precise locations within NWTT. Marine mammal and sea turtle dive behaviors are not easily stereotyped, but a species' behavior can generally be quantified by using an average percentage of time that an animal will typically spend within a range of depths, or between depth bins. For each species, a distribution throughout the water column is presented, along with a list of the references that are the sources of these data and an explanation of how these references were used to create the distribution. Depth bins are given in meters (m). Depending on the species, the distribution may cover a larger or smaller range of depths, such as for a shallow diving fur seal or a deep diving sperm whale. Likewise, depth bins may be smaller near the surface or larger at greater depths (e.g., 20 m bins near the surface where the animal spends more time or 100 m bins at the deepest depths the animal can reach).

2.2.1 Cetaceans

2.2.1.1 Family Balaenopteridae

2.2.1.1.1 *Balaenoptera acutorostrata*, Common Minke Whale

Minke whales are widely distributed throughout the world oceans, occurring in coastal and continental shelf waters, the deeper waters along continental slopes, and further seaward (Dorsey et al. 1990; Øien 1990). Fish (e.g., capelin, sandlance, and herring) and planktonic crustaceans (e.g., krill) are the main components of the minke whale diet (Haug et al. 1995). Minke whales feed by side-lunging into schools of prey as well as by gulping large amounts of water (Jefferson et al. 2008).

Little data have been collected on the dive behavior of minke whales. In order to build a representative depth distribution for minke whales, data from Figure 2 in Blix and Folkow (1995) were used. Blix and Folkow (1995) presented a time-depth record for a single minke whale tagged off the west coast of Svalbard, a Norwegian archipelago. This animal was predominantly foraging between 25 and 50 m. Two depth bins and the time spent within each depth bin were estimated, with the resulting depth distribution shown. The depth distribution data are derived from a short (75 minutes [min]) dive profile of a single animal, in which two behaviors are represented, cruising (52 percent of time) and foraging (48 percent of time); however, the amount of time spent in each of these two behaviors can vary significantly among individuals (Blix and Folkow 1995). The depth distribution for common minke whales is given in Table 2.

Table 2. Percentage of Time at Depth for the Minke Whale¹

Depth Bin (m)	% of Time at Depth
0–25	79.7
25–65	20.3

¹Based on data from Blix and Folkow (1995)

More recent data suggest that the common minke whale is capable of diving to greater depths than depicted by this distribution. For example, minke whales in the Antarctic have been associated with krill patches found at a median depth of 118 m (Friedlaender et al. 2009b). Off Scotland, minke whales are found where patches of pre-spawning herring occur at depths between 100 and 150 m (MacLeod et al. 2003b), while off the coast of California, tagged minke whales dove to 130 m (Southall et al. 2014), although in both cases whales spend the majority of time in the top 25 m of the water column. There is also limited evidence that minke whales may exhibit diurnal variation in diving behavior (Joyce et al. 1990; Stockin et al. 2001). The depth distribution shown in Table 2 will be considered representative for common minke whales until more information becomes available.

2.2.1.1.2 *Balaenoptera borealis*, Sei Whale

Sei whales have a cosmopolitan distribution, migrating between high-latitude feeding grounds and low-latitude breeding grounds (Horwood 2002). Sei whales are capable of diving for between 5 and 20 min (Reeves et al. 2002) to feed on plankton, predominantly copepods and euphausiids, which occur between the surface and depths around 150 m (Budylenko 1978; Flinn et al. 2002). They may also feed on small schooling fish and cephalopods by both gulping and skimming.

Little data have been collected on the dive behavior of sei whales. Sei whales are not thought to be deep divers. (Baumgartner et al. (2011) found that sei whales were absent during times when copepods were at depth, suggesting that sei whales may only be able to feed effectively on copepod aggregations when

they are at or near the surface. In addition, Baumgartner and Fratantoni (2008) observed low calling rates during the night when copepods were at the surface, and higher calling rates during the day when copepods were at depth. This study speculated that sei whales reduced calling rates to accommodate nighttime feeding on the copepod aggregations at the surface, and increased calling rates during the day when copepods migrated to deeper depths where they were unavailable as prey to the sei whales.

Due to a lack of available data on the dive behavior of sei whales, they will be represented by a surrogate species: the Bryde's whale (Section 2.2.1.1.3). The Bryde's whale is the closest relative to the sei whale (Sasaki et al. 2005); these species are of similar body size (Horwood 2002) and feed on similar prey in the Northern Hemisphere (Flinn et al. 2002; Mizroch et al. 1984). While sei whales differ from other Balaenopterids in their prey preference for copepods, this preference means that, like Bryde's whales, sei whales are not thought to be deep divers and spend most of their time near the surface (Alves et al. 2010). Foraging sei whales and Bryde's whales utilize similar water depths (Alves et al. 2010; Baumgartner et al. 2011). The depth distribution for the sei whale can be found in Table 3.

2.2.1.1.3 *Balaenoptera edeni*, Bryde's Whale

Bryde's whales are found in tropical and temperate waters, with separate coastal and offshore forms (Best 2001; Weir 2007). There is ongoing debate about the taxonomic relationship between two morphotypes — the larger *-brydei* form and the smaller *-edeni* form (Sasaki et al. 2006). These two forms are genetically distinct, and are differentiated by geographic distribution, inshore/offshore habitat preferences, and size. However, for both morphotypes, which are not easily distinguished at sea, the scientific name *B. edeni* is commonly used. The main prey of Bryde's whales include pelagic schooling fish species, such as sardines, mackerel, and herring (Siciliano et al. 2004), as well as cephalopods and small crustaceans (Kato 2002; Omura 1962).

In order to build a representative depth distribution for Bryde's whales, data from Table 1 in Alves et al. (2010) were used. Alves et al. (2010) reported a distribution of time spent in shallow versus deep dives for two whales tagged with a time-depth recorder near Madeira Island, Spain. Though these data are not strictly an indication of time spent in the two different depth bins (time spent diving to 40–292 m includes time passing through the 0–40 m depth bin), these data are the best available approximation of time spent at depth. The depth distribution for Bryde's whales is given in Table 3.

Table 3. Percentage Time at Depth for the Bryde's Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0–40	84.5
40–292	15.5

¹Based on data from Alves et al. (2010)

²This depth distribution is also representative of the sei whale

2.2.1.1.4 *Balaenoptera musculus*, Blue Whale

Blue whales have a cosmopolitan distribution, living in both coastal and offshore waters (Jefferson et al. 2008). Blue whales track the diel vertical migration of their prey and feed almost exclusively on euphausiids, or krill (Sears 2002). Although surface feeding has been observed during the daylight, it is more usual for blue whales to dive to at least 100 m into layers of euphausiid concentrations during daylight hours and feed nearer the surface at night (Sears and Perrin 2008).

In order to build a representative depth distribution for blue whales, data from Figures 4 and 8 in Oleson et al. (2007), as well as Figure 2 from Acevedo-Gutiérrez et al. (2002) were used. Oleson et al. (2007) provided graphs of the percent time at depth of 38 blue whales off the coast of California in Figure 8.

The data for the non-vocal, AB callers, and D callers were averaged together to get a general depth distribution. However, percentage of time at the surface was ignored by this study. By incorporating the average number of surfacing events over time in the sample dive profile from Figure 4 in Oleson et al. (2007) and the average time spent at surfacing events from Figure 2 in the Acevedo-Gutiérrez et al. (2002) study, a percent time spent in the surface bin could be estimated (20.9 percent). The remaining bins from the Oleson et al. (2007) study were redistributed proportionally to account for the remaining 79.1 percent of time. The depth distribution for blue whales is given in Table 4.

Table 4. Percentage of Time at Depth for the Blue Whale¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–5	20.9	155–165	2.2
5–15	12.7	165–175	2.3
15–25	9.9	175–185	2.2
25–35	6.4	185–195	2.2
35–45	5.3	195–205	1.4
45–55	4	205–215	1.1
55–65	3.5	215–225	1.1
65–75	2.8	225–235	1
75–85	2.2	235–245	0.9
85–95	2.4	245–255	0.7
95–105	2.4	255–265	0.7
105–115	2.1	265–275	0.3
115–125	2	275–285	0.2
125–135	2.1	285–295	0.2
135–145	2.1	295–305	0.2
145–155	2.3	305–315	0.2

¹Based on data from Oleson et al. (2007) and Acevedo-Gutiérrez et al. (2002)

While other studies did not include depth distributions for blue whales, they did provide additional information to categorize dive behavior. Blue whales in the Gulf of St. Lawrence conducted foraging dives to 150 m, where feeding lunges were observed (Doniol-Valcroze et al. 2011). Similarly, a study conducted in Monterey Bay found that blue whales fed on the most concentrated patches of krill at depths of 130 to 150 m (Schoenherr 1991). Blue whales off central California foraged at depths between 130 and 300 m (Calambokidis et al. 2008; Croll et al. 2001), while in southern California, dive depths ranged from 50 to 350 m (Acevedo-Gutiérrez et al. 2002; Croll et al. 2001; De Vos et al. 2012; Goldbogen et al. 2012; Goldbogen et al. 2013; Mate et al. 2016; Oleson et al. 2007; Southall et al. 2014). Seven whales tagged off the coast of southern California dove to a mean depth of 140 m and a maximum depth of 204 m during foraging, while to only a mean depth of 67.6 m during non-foraging dives (Croll et al. 2001). These data are consistent with the depth distribution in Table 4.

2.2.1.1.5 *Balaenoptera physalus*, Fin Whale

The fin whale occurs in greatest concentrations in cold and temperate waters around the globe and are commonly found seaward of the continental slope (Aguilar 2002). Prey species include euphausiids (Laidre et al. 2010; Ruchonnet et al. 2006; Vikingsson 1997), schooling fish such as herring and capelin (Nottestad et al. 2002), and cephalopods (Flinn et al. 2002). A 2001 study has shown that dense prey concentrations are typically found at depths greater than 100 m off the coast of California (Croll et al. 2001).

In order to build a representative depth distribution for fin whales, data from Figure 4a in Croll et al. (2001) and the text of Goldbogen et al. (2006) were used. Due to the lack of data on time spent at depth, the data from Croll et al. (2001) were used as a proxy for percentage of time spent at depth. Croll et al. (2001) found that, amongst the 15 tagged fin whales, there was a maximum dive depth of 316 m. Foraging dives were deeper and longer in duration than non-feeding dives. Goldbogen et al. (2006) reported that tagged whales spent 40 percent of time in the top 50 m. Time spent at depths deeper than 50 m were extracted from dive profiles presented in Croll et al. (2001) to represent the remaining 60 percent of time. The depth distribution for fin whales is given in Table 5.

Table 5. Percentage of Time Spent at Depth for the Fin Whale¹

Depth Bin (m)	% of Time at Depth
0–50	40
50–70	13.9
70–90	9.6
90–110	8.1
110–130	11.4
130–150	8.4
150–170	3.5
170–190	2.0
190–210	1.4
210–230	0.6
230–250	0.6
250–270	0.2
270–290	0.1
290–310	0.2

¹Based on data from Croll et al. (2001) and Goldbogen et al. (2006)

While other studies did not include depth distributions for fin whales, they did provide additional information to categorize dive behavior. Off southern California, foraging dives of 100 to 300 m (Acevedo-Gutiérrez et al. 2002; Goldbogen et al. 2006; Mate et al. 2016) have been recorded. In the Ligurian Sea, a maximum dive to over 470 m was noted (Panigada et al. 1999), while Southall et al. (2014) reported that dives by fin whales rarely exceeded 250 m. These data are consistent with the depth distribution in Table 5.

2.2.1.1.6 *Megaptera novaeangliae*, Humpback Whale

Humpback whales have a cosmopolitan distribution in the coastal and continental shelf waters of the globe. They migrate between mid to high-latitude foraging grounds and low-latitude breeding grounds (Clapham 2002). Humpback whales feed on a variety of organisms, including euphausiids and small schooling fish (Hain et al. 1982; Hazen et al. 2009; Laerm et al. 1997).

Due to a separation of behaviors based on location within a foraging ground or within a breeding ground, a separate representative depth distribution was compiled for humpback whales within foraging grounds. In order to build a representative depth distribution for humpback whales within foraging grounds, data from Figure 3.8 as well as the text of Dietz et al. (2002) were used. Dietz et al. (2002) shows the number of dives per hour to specific depth bins. The data from Dietz et al. (2002) were used as a proxy for percentage of time spent at depth for the six whales tagged off the foraging grounds of West Greenland. While Figure 3.8 begins at a depth of 8 m, the text states that the average time spent at the surface is 83.3 percent during mid-day and 75 percent at midnight, resulting in an average surface

time of 79.2 percent. The data from Dietz et al. (2002) were then redistributed proportionally to account for the remaining 20.8 percent of time. The depth distribution for humpback whales on foraging grounds is given in Table 6.

Table 6. Percentage of Time at Depth for Humpback Whales on Foraging Grounds¹

Depth Bin (m)	% of Time at Depth
0–8	79.2
8–20	11.5
20–35	2.7
35–50	1.0
50–100	1.6
100–150	1.4
150–200	1.2
200–300	1.2
300–400	0.2

¹Based on data from Dietz et al. (2002)

Dive depths on the Greenland foraging grounds are consistent with the depth of feeding reported by Goldbogen et al. (2008) off central California and Dolphin (1987b) off Alaska. Dolphin (1987a); however, reported that 75 percent of feeding dives were to less than 60 m, and Friedlaender et al. (2009a) found evidence of bottom feeding in the shallower water (less than 50 m) of the Gulf of Maine. While the depth distribution in Table 8 has a maximum depth of 400 m, over 95 percent of the time is in the top 50 m. Therefore, the depth distribution in Table 6 is consistent with these studies as well.

2.2.1.2 Family Delphinidae

2.2.1.2.1 *Delphinus delphis*, Short-beaked Common Dolphin

While species of common dolphins are sympatric in some nearshore continental shelf waters, short-beaked common dolphins are typically found in deeper waters along the continental slope (Cañadas and Hammond 2008; Heyning and Perrin 1994; Jefferson et al. 2009; Rosel et al. 1994; Selzer and Payne 1988). They feed on epipelagic and mesopelagic fish and squid (Selzer and Payne 1988), and they also forage at night on vertically migrating prey associated with the deep scattering layer (Evans 1994; Neumann and Orams 2003; Ohizumi et al. 1998; Pusineri et al. 2007).

Little data have been collected on the dive behavior of the short-beaked common dolphin. Evans (1975; 1994) described the late afternoon and evening diving behavior of an adult female short-beaked common dolphin in the Pacific Ocean. Before 1730 the dolphin mostly remained in the top 10 m, at which time it switched to a pattern of regular dives to 50 m, with a maximum dive depth of just over 200 m (Evans 1974, 1994).

Due to the lack of available data on the diving behavior of the short-beaked common dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.2.7). Pantropical spotted dolphins also make shallower dives during the day than at night, when they forage on vertically-migrating prey associated with the deep scattering layer (Scott and Chivers 2009). During the day, pantropical spotted dolphins spend 94 percent of their time in the top 20 m of the water column, while at night 95 percent of their time is spent in the top 50 m (Baird et al. 2001). Evans (1994) reported the maximum dive depth for three short-beaked common dolphins was 257 m. Similarly, maximum dive depths for pantropical spotted dolphins are 122 m for daytime and 213 m for nighttime (Baird et al. 2001). Pantropical spotted dolphins and common dolphins are members of the same subfamily,

Delphinidae (LeDuc et al. 1999), and their behavior shows clear similarities in diving pattern, foraging behavior, and water column usage. The depth distribution of the short-beaked common dolphin is given in Table 11.

2.2.1.2.2 *Globicephala macrorhynchus*, Short-finned Pilot Whale

Short-finned pilot whales occur in tropical and warm-temperate waters along the continental shelf and slope (Davis et al. 1998). Short-finned pilot whales feed predominantly on squid, but they also occasionally feed on octopus and fish (Mintzer et al. 2008; Reeves et al. 2002). On the U.S. Pacific coast, the neritic cephalopod, *Loligo* spp., is their dominant prey (Mintzer et al. 2008). Short-finned pilot whales feed on vertically migrating prey, diving deep during dusk and dawn and staying near surface at night (Baird et al. 2003).

In order to build a representative depth distribution for short-finned pilot whales, data from Figure 9 in Wells et al. (2013) were used. Wells et al. (2013) tagged two male pilot whales after a mass stranding event in the Florida Keys. While one of the individual tags stopped transmitting after 16 days, the other tag transmitted for a total of 67 days; thus, the representative depth distribution contains only the Wells et al. (2013) data from the individual with the longer transmission time. Due to the lack of data on time spent at depth, the proportion of dives made to specific depth ranges from Wells et al. (2013) will be used as a proxy for percentage of time spent at depth. The depth distribution for short-finned pilot whales is given in Table 7.

Table 7. Percentage of Time at Depth for the Short-Finned Pilot Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0–2	32.25
2–50	46.75
50–100	3.00
100–200	5.00
200–300	4.25
300–400	2.75
400–500	1.75
500–600	1.75
600–700	1.75
700–800	0.5
800–900	0.125
900–1,000	0.125

¹Based on data from Wells et al. (2013)

²This depth distribution is also representative of the following species:
pygmy sperm whale and dwarf sperm whale

Aguilar Soto et al. (2008) reported a maximum dive depth of 1,019 m for 23 whales near the Canary Islands, which is similar to the deepest depth bin in Table 7. While the maximum depth in Jensen et al. (2011) (roughly 700 m) is hundreds of meters shallower than in the distribution above, the total time spent deeper than this depth constitutes a very small percentage of the whale's total time.

2.2.1.2.3 *Grampus griseus*, Risso's Dolphin

Risso's dolphins are commonly found in temperate and tropical waters along continental slopes (Azzellino et al. 2008; Baumgartner 1997; Green et al. 1992). Although little is known about their foraging or diving behavior, vertically migrating cephalopods are presumed to be the primary food source for Risso's dolphins (Clarke and Pascoe 1985).

In order to build a representative depth distribution, data from Figure 5 in Wells et al. (2009) were used. Wells et al. (2009) reported on the movement and diving behavior of a rehabilitated adult male Risso's dolphin that stranded on the Gulf coast of Florida. Based on Figure 5, the depth distribution for Risso's dolphins was estimated for four 6-hour blocks of time. The tagged animal in this study travelled through waters with a mean depth of 548 m (range 3–2,300 m), and was therefore likely not diving close to the seafloor. The deepest dive recorded on the tag was in the 400–500 m depth range, and less than 0.1 percent of dives were deeper than 200 m (Wells et al. 2009). The average time spent in these depth bins was calculated for the representative depth distribution. The depth distribution for the Risso's dolphin is given in Table 8.

Table 8. Percentage of Time at Depth for the Risso's Dolphin¹

Depth Bin (m)	% of Time at Depth
0–1	24.8
1–2	13.5
2–10	16.5
10–50	43.5
50–100	1.2
100–150	0.1
150–600	0.4

¹Based on data from Wells et al. (2009)

2.2.1.2.4 *Orcinus orca*, Killer Whale

Killer whales have a cosmopolitan distribution, but they are most commonly observed in temperate, coastal waters (Ford 2002). Killer whales feed on a variety of prey, although most populations exhibit some degree of dietary specialization. In the northeastern Pacific and in the Antarctic, sympatric populations in each location are socially (and, in some cases, reproductively) isolated by foraging specializations for fish or marine mammal species (Ford et al. 1998; Pitman and Ensor 2003; Saulitis et al. 2000).

Due to a separation of diving behaviors based on preferred prey of the killer whale, two separate representative depth distributions were compiled for killer whales: fish-eating killer whales and mammal-eating killer whales. Fish-eating killer whales have been studied more extensively than mammal-eating ecotypes, although there is still limited published information on diving behavior of either. Fish-eating killer whales will either chase individual prey at the surface, or collectively herd schooling fish towards the surface (Domenici et al. 2000; Nøttestad et al. 2002). Mammal-eating killer whales have different foraging strategies than fish-eating killer whales (Barrett-Lennard et al. 1996; Pitman and Ensor 2003). Mammal-eating killer whales often attempt to capture prey from below, where a prey's silhouette against brighter surface waters may improve detection. Miller et al. (2010) found deeper dives for mammal-eating killer whales that occurred during the day.

In order to build a representative depth distribution for fish-eating killer whales, data from Figure 2 in Sivle et al. (2012), Figure 1e in Kvadsheim et al. (2012), as well as plots from post sonar exposure and/or silent pass of the ship from Miller et al. (2011) were used. Since all three studies analyzed the potential effects of sonar on the dive behavior of killer whales, only dive profiles from periods of time when no sonar was active were used. Depth distributions were extracted from the presented dive profiles. Individual depth distributions for each animal were averaged to create the representative depth distribution in Table 9. This representative depth distribution is consistent with Baird (1994) and Shapiro (2008), who reported that fish-eating resident killer whales spent the vast majority of their time in the top 20 m. The depth distribution for killer whales is given in Table 9. The deepest killer whale dive

recorded thus far was to 264 m by Baird et al. (2005a), who tagged a total of 34 Southern Resident killer whales. However, the average of all the tagged killer whale deepest dives was 141 m, which is consistent with the depth distribution in Table 9 below. This depth distribution will be used for the Northern Resident, Southern Resident, Alaska Resident, and East Pacific Offshore stocks of killer whales.

Table 9. Percentage of Time at Depth for Fish-Eating Killer Whales¹

Depth Bin (m)	% of Time at Depth
0–20	81.5
20–40	8.6
40–60	4.5
60–80	2.4
80–100	2.1
100–120	0.5
120–140	0.3
140–150	0.1

¹Based on data from Kvadsheim et al. (2012), Miller et al. (2011), and Sivle et al. (2012)

In order to build a representative depth distribution for mammal-eating killer whales (Table 10), data from Figure 5 in Miller et al. (2010) and Figure 2.5B from Baird (1994) were used. Based on visual inspection and interpretation of the Figures provided by Miller et al. (2010), an average dive distribution was created for the presented whales; visual inspection was also used to build a depth distribution from the Figure presented by Baird (1994). The calculated data from Miller et al. (2010) was weighted by a factor of 11 when creating the final dive profile for mammal-eating killer whales, due to the presence of 11 animals in that study compared to one presented by Baird (1994). This depth distribution will be used for West Coast Transient killer whales.

Table 10. Percentage of Time at Depth for Mammal-Eating Killer Whales¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	42.9	50–60	2.5
10–20	23.4	60–70	1.7
20–30	14.0	70–80	0.8
30–40	8.3	80–100	0.4
40–50	6.0		

¹Based on data from Baird (1994) and Miller et al. (2010)

2.2.1.2.5 *Lagenorhynchus obliquidens*, Pacific White-sided Dolphin

Pacific white-sided dolphins inhabit cold temperate waters of the North Pacific, in both offshore and coastal waters (Brownell et al. 1999; Waerebeek and Würsig 2002). Their primary prey species include mesopelagic fish and cephalopods, as well as epipelagic fish in shallower waters (Brownell et al. 1999; Kajimura and Loughlin 1988; Miyazaki et al. 1991; Morton 2000; Walker and Jones 1994).

Little data have been collected on the dive behavior of the Pacific white-sided dolphin. Hall (1970) trained a captive Pacific white-sided dolphin to dive to a depth of 214 m. However, Black (1994) reported that in coastal waters, 70 percent of dives were shorter than 20 seconds in duration, and dives longer than 90 seconds were rare, indicating that most dives are shallow. Heise (1997) similarly reported that 70 percent of foraging dives were less than 15 seconds in duration. Therefore, Pacific white-sided dolphins are not considered deep divers. This species is thought to feed mostly at night or in the morning (Stroud et al. 1981) when their mesopelagic prey rise to surface waters.

Due to the lack of available data on the diving behavior of the Pacific white-sided dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.2.7). Pantropical spotted dolphins spend the majority of their time in the top 50 m, and their maximum diving depths are within the range of the dive depth of the trained Pacific white-sided dolphin (Baird et al. 2001; Hall 1970; Scott and Chivers 2009). Pantropical spotted dolphins and Pacific white-sided dolphins also feed on similarly migrating prey species. The depth distribution for the Pacific white-sided dolphin can be found in Table 11.

2.2.1.2.6 *Lissodelphis borealis*, Northern Right Whale Dolphin

The northern right whale dolphin is abundant in deep, temperate waters across the North Pacific (Forney and Barlow 1998; Jefferson and Newcomer 1993b; Leatherwood and Walker 1979; Rankin et al. 2007). They are known to commonly associate with Pacific white-sided dolphins and Risso's dolphins (Forney and Barlow 1998; Jefferson and Newcomer 1993b), with which they show dietary overlap (Walker and Jones 1994). Northern right whale dolphins near the southern California coast feed principally on cephalopods and a diverse variety of myctophid fish (Jefferson and Newcomer 1993a; Jefferson et al. 1994; Leatherwood and Walker 1979).

Little data have been collected on the dive behavior of the northern right whale dolphin. Some evidence based on stomach contents suggests that northern right whale dolphins may dive as deep as 200 m (Fitch and Brownell 1968; Jefferson et al. 1994). Individual northern right whale dolphins have been observed to dive for brief periods (10 to 75 seconds), but can also remain submerged for up to 6.5 min (Cruikshank and Brown 1981; Leatherwood and Walker 1979). Northern right whale dolphins have comparatively low muscle myoglobin content among odontocetes, suggesting they are not deep divers (Noren and Williams 2000).

Due to the lack of available data on the diving behavior of the northern right whale dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.2.7). Of the two dolphins with which the northern right whale dolphin associates the most, the Risso's dolphin is considered a much deeper diver than the Pacific white-sided dolphin. Because northern right whale dolphins have low muscle myoglobin content and are thought to feed on prey only as deep as 200 m, they are considered shallower divers. Therefore, due to dietary similarity and frequent association with the Pacific white-sided dolphin, the northern right whale dolphin will be represented by the same surrogate species, the pantropical spotted dolphin. The depth distribution for the Northern right whale dolphin can be found in Table 11.

2.2.1.2.7 *Stenella attenuata*, Pantropical Spotted Dolphin

Pantropical spotted dolphins are found in warm-temperate and tropical waters over the continental slope and offshore in deeper waters (Perrin and Hohn 1994). Pantropical spotted dolphins feed on both epipelagic and mesopelagic fish and squid (Wang et al. 2003). In general, pantropical spotted dolphins dive deeper at night, foraging on prey associated with vertical migrations of the deep scattering layer (Robertson and Chivers 1997; Scott and Chivers 2009).

In order to build a representative depth distribution for pantropical spotted dolphins, data from Figure 4 and Table 2 in Baird et al. (2001) and Figure 9 and Table 2 in Scott and Chivers (2009) were used. While Baird et al. (2001) looked at pantropical spotted dolphin diving behavior around the Hawaiian Islands, Scott and Chivers (2009) recorded data on these dolphins in pelagic waters. Baird et al. (2001) reported pantropical spotted dolphins spend, on average, 88.5 percent of their time within 10 m of the surface during the day. Baird et al. (2001) reported the daytime average percentage of time in two meter intervals for the top 10 meters. For the Baird et al. (2001) nighttime and Scott and Chivers (2009) data,

the percentage of time determined for the top 10 meters was uniformly distributed across these two meter intervals. Daytime and nighttime averages were calculated for the Baird et al. (2001) data, and these were then averaged with the Scott and Chivers (2009) data. The resulting mean daytime and nighttime depth distribution data are presented in Table 11. Baird reported maximum daytime and nighttime dive depths at 122 m and 213 m, respectively (Baird et al. 2001); however, Scott and Chivers (2009) calculated that dives to more than 120 m accounted for less than 0.1 percent of all dives. They also noted that daytime dives were primarily shallow and above the thermocline (Scott and Chivers 2009). The depth distribution for pantropical spotted dolphins is given in Table 11.

Table 11. Percentage of Time at Depth for the Pantropical Spotted Dolphin^{1,2}

Depth Bin (m)	% of Time at Depth	DeDpth Bin (m)	% of Time at Depth
0–2	20.3	70–80	0.6
2–4	10.6	80–90	0.6
4–6	8.6	90–100	0.4
6–8	9.0	100–110	0.3
8–10	9.5	110–120	0.3
10–20	21.2	120–130	0.1
20–30	8.8	130–140	0.1
30–40	3.8	140–150	0.1
40–50	2.5	150–160	0.1
50–60	1.9	160–170	0.1
60–70	1.1		

¹Based on data from Baird et al. (2001) and Scott and Chivers (2009)

²This depth distribution is also representative of the following species: the long-beaked common dolphin, short-beaked common dolphin, Pacific white-sided dolphin, Northern right whale dolphin, and striped dolphin

2.2.1.2.8 *Stenella coeruleoalba*, Striped Dolphin

Striped dolphins prefer tropical and warm-temperate waters and have an oceanic distribution, with most observations occurring beyond the continental shelf (Archer II 2002; Cañadas et al. 2002; Davis and Fargion 1996; Davis et al. 1998; Perrin et al. 1994). Striped dolphins primarily feed on small, pelagic, vertically migrating prey (Blanco et al. 1995). Stomach contents analyses suggest that foraging occurs mostly in the dusk and early evening hours (Ringelstein et al. 2006). Their distribution in the North Atlantic Ocean is associated with a mesopelagic prey community comprised of fish and cephalopod species (Doksaeter et al. 2008).

Little data have been collected on the dive behavior of the striped dolphin. A single striped dolphin carrying a time-depth recorder dove to a mean depth of 22.6 m standard deviation (SD) =17.5 during the day and 126.7 m (SD=120.9) at night, with a maximum dive depth of 705 m (Minamikawa et al. 2003).

Due to the lack of available data on the diving behavior of the striped dolphin, it will be represented by a surrogate species: the pantropical spotted dolphin (Section 2.2.1.2.7). The observed pattern of shallow daytime diving and deeper nighttime diving reported in Minamikawa et al. (2003) is consistent with similar diving behavior seen in short-beaked common dolphins (Section 2.2.1.2.1) and their surrogate species, pantropical spotted dolphins (Section 2.2.1.2.7), which are also in the genus *Stenella*. Additionally, all three species occur in similar water depths (Davis et al. 1998). However, it is acknowledged that the striped dolphin may dive to deeper depths on average, due to the deep maximum dive depth recorded by Minamikawa et al. (2003). The depth distribution for striped dolphins is given in Table 11.

2.2.1.2.9 *Tursiops truncatus*, Bottlenose Dolphin

Bottlenose dolphins have a cosmopolitan distribution in the tropical and temperate waters of the world (Wells and Scott 2002). They reside in estuarine, coastal, and offshore continental shelf and slope waters. Populations vary in their migratory and foraging behavior (Wells and Scott 2002). Bottlenose dolphins feed primarily on fish species, with squid and other invertebrates contributing to their diet as well (National Marine Fisheries Service 2015). Due to the range of habitats in which bottlenose dolphins are found, prey species may be epipelagic, pelagic, mesopelagic, or benthic in origin, depending on the region and habitat (Mead and Potter 1990; Rossbach and Herzing 1997; Shane 1990; Wells and Scott 1999). The presence of deep-sea fish in the stomachs of some offshore animals suggests that they can dive to depths greater than 500 m (Reeves et al. 2002).

Little data have been collected on the dive behavior of bottlenose dolphins. In order to build a representative depth distribution for bottlenose dolphins, data from Table 4 in Klatsky (2004) were used. Dolphins 39999 and 40000 were tagged for 30 and 48 hours, respectively, whereas Dolphin 40001 was tracked for 45 days, providing 792 hours of dive data. So, while Klatsky (2004) presents the percentage of time at depth for three individuals, only the data from Tag 40001 were considered for the depth distribution in order to provide the most comprehensive view of bottlenose dolphin diving behavior. Klatsky (2004) reported that the maximum recorded depth of a dolphin was 492 m; therefore, that value was used as the maximum depth associated with this depth distribution. The depth distribution for bottlenose dolphins is given in Table 12.

Table 12. Percentage of Time at Depth for the Bottlenose Dolphin¹

Depth Bin (m)	% of Time at Depth
0–6	64.6
6–10	3.9
10–26	5.7
26–50	2.5
50–76	2.0
76–100	1.7
100–150	3.6
150–200	2.5
200–250	2.1
250–300	2.2
300–350	1.9
350–400	1.7
400–450	1.6
400–492	4.0

¹Based on data from Klatsky et al. (2004)

These data are consistent with animals foraging up to 500 m off Hawaii and spending the majority of their time between the surface and 50 m (Baird et al. 2014).

2.2.1.3 Family Eschrichtiidae

2.2.1.3.1 *Eschrichtius robustus*, Gray Whale

Gray whales are distributed coastally throughout the Pacific Ocean, migrating annually between Arctic and subtropical waters (Jones and Swartz 2002; Swartz 1986). Gray whales forage within the water column, with modified skimming techniques used to capture neritic fish, and scraping along the benthos

to acquire benthic fish, squid, annelids, crustaceans, and mollusks (Darling et al. 1998; Dunham and Duffus 2002; Jones and Swartz 2002; Nerini 1984). Gray whales have been reported foraging in water up to 120 m deep (Cacchione et al. 1987; Dunham and Duffus 2002; Würsig et al. 1986), although whales forage in waters less than 20 m deep in many areas (Guerrero 1989; Ljungblad et al. 1987; Malcolm and Duffus 2000; Malcolm et al. 1995; Stewart et al. 2001; Woodward and Winn 2006).

In order to build a representative depth distribution for gray whales, data from Figure 3 in Malcolm et al. (1995) were used. Malcolm et al. (1995) reported the percentage of time at depth for a single foraging whale carrying a tag for over 8 hours in waters off British Columbia in an area with a mean bottom depth of 18 m. The majority of dives (76 percent) were ventilation dives (to a mean depth of 2.3 m), while 13 percent were feeding dives (to a mean depth of 16.7 m). The whale appeared to spend little time at intermediate depths within the water column, spending most of its time either breathing at the surface or feeding at the bottom. Due to the large size of gray whales, the data from Malcolm et al. (1995) was summed into four-meter bins for the representative depth distribution. The depth distribution for gray whales is given in Table 13.

Table 13. Percentage of Time at Depth for the Gray Whale¹

Depth Bin (m)	% of Time at Depth
0–4	39.0
4–8	8.5
8–12	7.0
12–16	16.0
16–20	28.0
20–22	1.5

¹Based on data from Malcolm et al. (1995)

While other studies did not include depth distributions for gray whales, they did provide additional information to categorize dive behavior. The representative depth distribution data compare to a later study with a larger sample size of whales, where 79 percent of dives by whales off Vancouver Island were to a mean depth of 2.2 m, and 15 percent of dives were to a mean depth of 12–19 m (Malcolm and Duffus 2000). Woodward and Winn (2006) and Woodward (2006) similarly reported that six whales feeding along the central British Columbia coast had a mean dive depth of 11 m (range 2.4–28.9 m). The percentage of time near the surface (19.5 percent from 0–2 m) is also consistent with other studies in the same region (14.2 percent and 17.5 percent) (Stelle et al. 2008) and in the Bering Sea (22 percent) (Würsig et al. 1986). Furthermore, the dive depth is similar to the reported foraging depths in British Columbia and other regions (Guerrero 1989; Ljungblad et al. 1987; Malcolm and Duffus 2000; Malcolm et al. 1995; Stewart et al. 2001; Woodward and Winn 2006). Stewart et al. (2001) described the diving behavior post-release for a rehabilitated calf in southern California; all dives were less than 20 m deep, and 85 percent of dives were less than 10 m deep. An earlier release of a post-rehabilitated calf in the same area documented a much deeper maximum diving depth (170 m) and an average diving depth of approximately 50 m (Evans 1974), which is deeper than the that in the representative depth distribution.

2.2.1.4 Family Kogiidae

2.2.1.4.1 *Kogia spp.* Guild

Pygmy (*Kogia breviceps*) and dwarf (*Kogia sima*) sperm whales have a cosmopolitan distribution in all temperate and tropical waters (Bloodworth and Odell 2008). Little data have been collected on the dive

behavior of either species. Sightings of pygmy sperm whales in the North Atlantic are most common in waters ranging from 400 to 1,000 m in depth (Clarke 2003; Scott et al. 2001; Waring et al. 2006). Mid- and deep-water cephalopods predominantly contribute to the diet of the pygmy sperm whale (Beatson 2007; Bloodworth and Odell 2008; Fernandez et al. 2009; McAlpine et al. 1997; Ross 1979; Santos et al. 2006). Based on the analysis of the stomach contents of whales stranded in New Zealand, Beatson (2007) concluded that pygmy sperm whales feed at shallower depths within the water column than sperm whales, although some prey species are found at depths greater than 600 m. Similarly, Ploen (2004) found that prey species from the stomachs of stranded pygmy sperm whales in South Africa are found at depths below 300 m. There is some indication that dwarf sperm whales have a more coastal distribution than pygmy sperm whales, and prey often include more continental shelf and slope species than those of the pygmy sperm whale (Ross 1979; Wang et al. 2002).

Within the NWTT Study Area, the two *Kogia* species are difficult to distinguish when viewed at sea and are found in overlapping areas. Therefore, much of the density data from surveys is combined for both species into a generic *Kogia* spp. guild. Due to the lack of available data on the diving behavior of the pygmy or dwarf sperm whale, the two species will be represented by the surrogate species: the short-finned pilot whale (Section 2.2.1.2.2). The short-finned pilot whale is another primarily squid-eating species (Mintzer et al. 2008; Reeves et al. 2002), which forages deep in the water column (Jensen et al. 2011). The broad similarity in prey types and oceanic habitat suggests similarity in diving behavior to the short-finned pilot whale. The depth distribution for *Kogia* spp. can be found in Table 7.

2.2.1.5 Family Phocoenidae

2.2.1.5.1 *Phocoena phocoena*, Harbor Porpoise

Harbor porpoises inhabit temperate and sub-arctic continental shelf waters in the northern hemisphere. Their diet consists primarily of fish, including both pelagic schooling and benthic species (Bjorge and Tolley 2002; Recchia and Read 1989). Cephalopods, crustaceans, euphausiids, and polychaetes also contribute to their overall diet (Recchia and Read 1989; Smith and Read 1992; Walker et al. 1998).

In order to build a representative depth distribution for harbor porpoises, data from Figure 3 in Otani et al. (1998); Figures 2 and 3 in Westgate et al. (1995); Figure 1 in Otani et al. (2000); Figures 2, 3, and 4 in Cooper et al. (1993); and Figure 7 in Westgate and Read (1998) were used. Cooper et al. (1993) reported that porpoises in the Bay of Fundy were capable of diving to 150 m but spent most of their time in the top 50 m of the water column. Westgate et al. (1995) reported porpoises diving to a maximum of 226 m, but that average dive depth for individual porpoises ranged from 14–41 m, and the depth range with the greatest proportion of dives was 2–10 m. Otani et al. (1998) found that harbor porpoises off the coast of Japan spent 74–86 percent of their time in the top 20 m of the water column, with an average dive depth of 12–19 m. The time at depth was visually inspected and averaged from all of the above figures to create the depth distribution for the harbor porpoise. This was done to include a total of 14 different harbor porpoises into the dive distribution. While data from Cooper (1993), Otani (1998), and Westgate et al. (1995) show the number or frequency of dives to specific depth bins, this data will be used as a proxy for percentage of time spent at depth. The depth distribution for harbor porpoises is given in Table 14.

Table 14. Percentage of Time at Depth for the Harbor Porpoise¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	39	110–120	0.4
10–20	17.8	120–130	0.4
20–30	12.7	130–140	0.3
30–40	10.1	140–150	0.2
40–50	6.9	150–160	0.2
50–60	4.6	160–170	0.1
60–70	2.5	170–180	0.1
70–80	1.5	180–190	0.1
80–90	1.4	190–200	0.1
90–100	1	200–210	0.1
100–110	0.4	210–226	0.1

¹Based on data from Cooper (1993), Otani et al. (2000), Otani et al. (1998), Westgate and Read (1998), and Westgate et al. (1995)

While other studies did not include depth distributions for harbor porpoises, they did provide additional information to categorize dive behavior. In Linnenschmidt et al. (2013), three harbor porpoises were tagged; two of the harbor porpoises showed consistent diving activity throughout the day, while one harbor porpoise showed a diel diving pattern with few dives during the day. Teilmann et al. (2013) found that harbor porpoises spent more time between 0 to 2 m at night than during the day; this may be due to the movement of their prey throughout the water column (e.g., herring [*Clupea harengus*] and sprat [*Sprattus sprattus*]) (Cardinale et al. 2003).

2.2.1.5.2 *Phocoenoides dalli*, Dall's Porpoise

Dall's porpoises can be found in the subarctic and cool temperate waters of the North Pacific Ocean, including the Bering Sea, Okhotsk Sea, and Sea of Japan (Jefferson 2002). Primary prey species include epipelagic and mesopelagic schooling fish and cephalopod species (Jefferson 1988; Ohizumi et al. 2000; Stroud et al. 1981; Walker 1996).

In order to build a representative depth distribution for the Dall's porpoise, data from Figures 3 and 4 in Baird and Hanson (1998) were used. Baird and Hanson (1998) tagged three Dall's porpoises with time-depth recorders in the waters between Washington State and British Columbia. Each animal had a median dive depth of less than 40 m, and maximum dive depths ranged from 197–278 m. Data from the tagged Dall's porpoises were averaged together to create the representative depth distribution in Table 15.

Table 15. Percentage of Time at Depth for the Dall's Porpoise¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–1	5.3	50–60	4.8
1–2	15.7	60–70	4.7
2–3	8.8	70–80	3.8
3–4	3.2	80–90	2.3
4–5	3.8	90–100	2
5–6	1.7	100–110	1.1
6–7	1	110–120	0.7
7–8	1.3	120–130	0.4
8–9	1.3	130–140	0.4
9–10	1.7	140–150	0.4
10–11	1.3	150–160	0.3
11–20	11.7	160–170	0.3
20–30	8.3	170–180	0.3
30–40	6.8	180–190	0.3
40–50	6.2	190–200	0.1

¹Based on data from Baird and Hanson (1998)

The representative depth distribution is consistent with stomach contents analyses which suggest that Dall's porpoises feed high in the water column on vertically migrating mesopelagic species but occasionally forage on deeper benthic prey (Jefferson 1988; Ohizumi et al. 1998).

2.2.1.6 Family Physeteridae

2.2.1.6.1 *Physeter macrocephalus*, Sperm Whale

The sperm whale has a cosmopolitan distribution, preferring deeper waters seaward of the continental shelf edge (Whitehead 2002). Females and immature males tend to inhabit tropical and temperate waters below 40 degrees North (°N) latitude, while maturing and adult males move to higher latitudes, occurring in polar waters as adults (Whitehead 2002). Sperm whales feed on cephalopod species, primarily squid, as well as mesopelagic and demersal fish and occasionally crustaceans (Fiscus et al. 1989; Flinn et al. 2002; Kawakami 1980; Martin and Clarke 1986).

To account for published differences in the foraging dive behavior of whales in different regions, separate depth distributions were generated for the Atlantic Ocean, Gulf of Mexico, and the Pacific Ocean. In general, time spent at depth for the regions is consistent with foraging dives to 800–1,200 m in the Western North Atlantic Ocean (Sivle et al. 2012; Teloni et al. 2008; Watwood et al. 2006), and 400–1,300 m in the Western North Pacific Ocean (Amano and Yoshioka 2003; Aoki et al. 2012; Aoki et al. 2007). Overall, sperm whales typically spend 70–80 percent of their time between 20 and 400 m (Sivle et al. 2012; Teloni et al. 2008). At mid- and low-latitudes, females and immature animals undertake stereotypic dives lasting about 45 min and to depths between 400 and 1,200 m (Teloni et al. 2008; Watwood et al. 2006). Off Japan, females and immature sperm whales performed similarly stereotyped dive patterns to 1,400 m, lasting 30–50 min (Aoki et al. 2012). Radically different dive behavior has been observed at high latitudes, where mature males undertake dives lasting up to 60 min and to depths of nearly 1,900 m (Sivle et al. 2012; Teloni et al. 2008).

To build a representative depth distribution for sperm whales in the Pacific Ocean, data from Aoki et al. (2007) were used. Aoki et al. (2007) tagged four whales off the coast of Japan. The mean dive depth for

nighttime was 515 m averaged over the two tag locations; the mean dive depth for the daytime was 749.5 m. While this may suggest a diel diving pattern that follows the availability of prey, the pattern seems to depend largely on location (Aoki et al. 2012). The depth distribution for sperm whales in the Pacific Ocean is given in Table 16.

Table 16. Percentage of Time at Depth for the Sperm Whale in the Pacific Ocean¹

Depth Bin (m)	% of Time at Depth
1–50	30.2
51–100	4.8
101–150	3.3
151–200	3.3
201–250	3.1
251–300	2.5
301–350	2.5
351–400	3.6
401–450	5.3
451–500	6.5
501–600	9.8
601–700	6.5
701–800	8.6
801–900	8.4
901–1000	1.3
1001–1100	0.3

¹Based on data from Aoki et al. (2012)

2.2.1.7 Family Ziphiidae

2.2.1.7.1 *Berardius bairdii*, Baird's Beaked Whale

Baird's beaked whales inhabit temperate waters of the North Pacific Ocean and adjoining seas, primarily in the deep waters offshore of the continental shelf (Balcomb 1989; Kasuya 1986). This species consumes benthic and epibenthic fish and cephalopods, and occasionally feeds on mesopelagic species as well (Balcomb 1989; Kasuya 2002; Walker et al. 2002).

Little data have been collected on the dive behavior of Baird's beaked whales. Stomach contents analysis suggests that whales are feeding at depths of 800–1,200 m off Japan, feeding on prey at or near the seafloor (Reeves et al. 2002; Walker et al. 2002). Minamaka (2007) reported that one animal carrying a time-depth recorder dove down to a maximum depth 1,777 m, with dives lasting up to 64.4 min, which is similar to the maximum dive duration of 67 min observed by Kasuya (1986). The maximum dive depth reported by the first deployment of a multi-sensor tag on this species was given as roughly 1,400 m (Stimpert et al. 2014).

Due to the lack of data on the diving behavior of the Baird's beaked whale, it will be represented by a surrogate species: Cuvier's beaked whale (Section 2.2.1.7.3). The Cuvier's beaked whale is also a member of the subfamily Ziphiidae, and the feeding habits and types of prey for the two species are similar. The diving pattern of Baird's beaked whales appears very similar to other beaked whales, in which a long duration, deep dive is followed by shorter duration, shallow dives (Minamikawa et al. 2007; Tyack et al. 2006). The depth distribution for the Baird's beaked whale can be found in Table 18.

2.2.1.7.2 *Mesoplodon densirostris*, Blainville's Beaked Whale

Blainville's beaked whales inhabit deep temperate and tropical waters of the world's oceans (Pitman 2002). Little is known about prey species, but the diet of Blainville's beaked whales includes mesopelagic cephalopods, fish, and crustaceans (Herman et al. 1994; Hickmott 2005; MacLeod et al. 2003a; Mead 1989).

In order to build a representative depth distribution for Blainville's beaked whales, data were acquired from Figures 3a and 3b from Arranz et al. (2011), Figure 6 from Baird et al. (2005b), Figures 3a and 3b from Baird et al. (2006), Figure 1 from Barlow et al. (2013), Digital Acoustic Recording Tag (DTAG) data from Johnson and Aguilar de Soto (2008b) and Johnson and Aguilar de Soto (2008a), DTAG data from Tyack (2010), and Figure 1b from Tyack et al. (2006). Arranz et al (2011) tagged 9 whales to collect acoustic and movement data, looking to study buzz and click behaviors during dives; Figures 3a and 3b show the dive profile of a male Blainville's beaked whale over a period of 17 hours. Baird et al. (2005b) tagged four Blainville's beaked whales and presented cumulative percentage of time spent at depth for two individuals: an adult female with young calf, the daytime data for a large sub-adult or adult female, and the nighttime data for the same sub-adult or adult female. Different data from that same female whale were used to create another set of dive profiles after a 22.6 hour deployment, as published in Baird et al. (2006). Barlow et al. (2013) used DTAGs to collect acoustic information for beaked whales in an attempt to estimate density and abundance; Figure 1 is a typical dive profile for a tagged whale captured over a 15-hour period. Raw DTAG data for two animals in the Canary Islands, provided by Johnson and Aguilar de Soto (Johnson and Aguilar de Soto 2008a, 2008b), were binned, as well as raw DTAG data from one animal in the Bahamas by Tyack (2010). Tyack et al. (2006) used DTAGs to create a representative dive profile for Blainville's beaked whale in an attempt to study how depth impacts foraging tactics. Data from each source were arranged into 100 m bins, and those bins were averaged together to create a representative depth distribution. The depth distribution for Blainville's beaked whales is given in Table 17.

Table 17. Percentage of Time at Depth for the Blainville's Beaked Whale^{1,2}

Depth Bin (m)	% of Time at Depth
1–100	54.3
100–200	10.2
200–300	3.7
300–400	3.2
400–500	3.7
500–600	3.4
600–700	3.7
700–800	4.1
800–900	4.7
900–1,000	3
1,000–1,100	2.2
1,100–1,200	1.8
1,200–1,300	1.1
1,300–1,400	0.8
1,400–1,500	0.1

¹Based on data from Arranz et al. (2011), Baird et al. (2005b), Baird et al. (2006), Barlow et al. (2013), Johnson and Aguilar de Soto (Johnson and Aguilar de Soto 2008a, 2008b), Tyack et al. (2006), and Tyack (2010)

² This depth distribution is also representative of the NWTT beaked whale guild

2.2.1.7.3 *Ziphius cavirostris*, Cuvier's Beaked Whale

Cuvier's beaked whales inhabit slope waters with steep gradients around the world's oceans, with the exception of the high polar seas (Heyning 1989). Stomach contents analyses indicate that prey species include mesopelagic and benthic cephalopods, fish, and crustaceans (Heyning 1989; Hickmott 2005; Santos et al. 2001). It appears, however, that Cuvier's beaked whales eat mostly squid, and the majority of prey are open-ocean species that occur well below the surface, including on or near the seafloor in deep waters (Reeves et al. 2002).

In order to build a representative depth distribution for the Cuvier's beaked whale (Table 18), data were acquired from Figure 1 from Aguilar de Soto et al. (2006), Figure 5 from Baird et al. (2005b), Figure 3c from Baird et al. (2006), Figure 2a from Baird et al. (2008), Figure 1 from Barlow et al. (2013), DTAG data from Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), Figure 1a from Kvadsheim et al. (2012), and Figures 2a and 3a from Schorr et al. (2014). Aguilar de Soto et al. (2006) presented a time-depth profile of a Cuvier's beaked whale off Italy over a 15.6 hour period. Baird et al. (2005b) presented the cumulative percentage of time spent at depth for an adult female during the day and at night. Similarly, Baird et al. (2008) looked at diel variation in Cuvier's beaked whale diving behavior, presenting the cumulative percentage of time spent at depth for two tagged whales during both the day and night. Barlow et al. (Barlow et al. 2013) used DTAGs to collect acoustic information for beaked whales in an attempt to estimate density and abundance; Figure 1 is a typical dive profile for a tagged whale captured over a 15-hour period. Raw DTAG data collected on multiple occasions for an animal in Liguria, Italy, provided by Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), were binned. Kvadsheim et al. (2012) presented changes in dive behavior in response to sonar; as this report is portraying normal Cuvier's beaked whale behavior, the portion of the dive profile provided by Kvadsheim et al. (2012), in which sonar was used, has been omitted from the calculated typical depth distribution. Schorr et al. (2014) provided multi-day tag data for Cuvier's beaked whales, allowing for observation of diel patterns in dive behavior. The depth distributions from these studies were averaged together to create the

representative depth distribution below. The depth distribution for Cuvier's beaked whales is given in Table 18.

Table 18. Percentage of Time at Depth for Cuvier's Beaked Whale^{1,2}

Depth Bin (m)	% of Time at Depth
0–100	32
100–200	10.3
200–300	11.7
300–400	5.6
400–500	4.1
500–600	3.9
600–700	5.1
700–800	4.1
800–900	3.5
900–1,000	4.8
1,000–1,100	4.4
1,100–1,200	3.3
1,200–1,300	2.2
1,300–1,400	1.4
1,400–1,500	1.1
1,500–1,600	0.3
1,600–1,700	0.9
1,700–1,800	0.5
1,800–1,900	0.8

¹Based on data from Aguilar de Soto et al. (2006), Baird et al. (2005b), Baird et al. (2006), Baird et al. (2008), Barlow et al. (2013), Johnson and Sturlese (Johnson and Sturlese 2008a, 2008b), Kvadsheim et al. (2012), and Schorr et al. (2014)

²This depth distribution is also representative of Baird's beaked whale

This representative depth distribution is consistent with foraging dives from 689 to 1,888 m in the Mediterranean Sea (Tyack et al. 2006) and to 1,450 m off Hawaii (Baird et al. 2006). Based on the representative depth distribution in Table 18, Cuvier's beaked whales spent 31.9 percent of their time between 0–100 m and 36.1 percent of time deeper than 500 m; these values remain consistent with the data reported by Baird et al. (2008), in which Cuvier's beaked whales spend 12.4–51.1 percent of their time spent at depths less than 50 m, and between 33.9–52.1 percent of time at depths greater than 500 m.

2.2.1.7.4 NWTT Small Beaked Whale Guild

Within the NWTT Study Area, all beaked whales are represented as one group due to the difficulty of identifying beaked whales at sea to the individual species level. The NWTT beaked whale guild is a composite of all species that would be present in the Study Area, including Cuvier's beaked whale (*Ziphius cavirostris*), Blainville's beaked whale, ginkgo-toothed beaked whale (*Mesoplodon ginkgodens*), lesser beaked whale (*Mesoplodon peruvianus*), Hubb's beaked whale (*Mesoplodon carlhubbsi*), Perrin's beaked whale (*Mesoplodon perrini*), and Stejneger's beaked whale (*Mesoplodon stejnegeri*). While both Blainville's and Cuvier's beaked whales are located within the NWTT Study Area, the depth distribution for Blainville's beaked whales (Section 2.2.1.7.1) was used as the representative species for this group. Blainville's beaked whales spend a greater percentage of time in the upper 100 m of water than

Cuvier's, making the model a more conservative estimate of affected beaked whales. The depth distribution of beaked whales in the NWTT study area can be found in Table 17.

2.2.2 Carnivores

2.2.2.1 Family Mustelidae

2.2.2.1.1 *Enhydra lutris*, Northern Sea Otter

The sea otter is a species that ranges around the North Pacific Ocean rim, from Baja California, Mexico to the east coast of the Russian Kamchatka peninsula and the Kuril Islands towards Japan. Sea otters inhabit nearshore waters. Typically, due to water depths, foraging would occur closer to shore and resting may occur nearshore or further offshore (Laidre et al. 2009). Off the coast of Washington, Laidre et al. (2009) found that adult females spent 60 percent of their time foraging in habitats no more than 10 m deep and were rarely found foraging in waters 30 m deep. Males in the same study spent most of their time (32-34 percent) foraging in habitats 10-30 m deep. Beyond 40 m, foraging was minimal (1-2 percent of time) for both sexes (Laidre et al. 2009). Sea otter prey includes benthic invertebrates (abalone, sea urchins, mussels, clams, snails, crabs, and worms) and occasionally bottom fish (Kenyon 1981; Laidre and Jameson 2006; Riedman and Estes 1990). Off the coast of Washington, females tend to rest 1,000 m offshore (42 percent of their resting time) and were rarely observed resting beyond 2,000 m offshore (Laidre et al. 2009). Males tended to rest 1,000-1,500 m offshore 40 percent of the time and 2,000-2,500 m offshore 11 percent of the time (Laidre et al. 2009).

Early studies of sea otter dive behavior assumed average dives to be within 10-30 m of the surface (Kenyon 1981), but one record reported recovery of a sea otter carcass from a crab pot set at 97 m (Newby 1975), and assumed the otter dove to that depth, but perished after getting stuck in the crab pot. Thometz et al. (2016) examined the foraging dive behavior of southern sea otters off the coast of California and found that the deepest foraging dives were performed by males (the maximum depth was 88 m), while the maximum dive by a female was 69 m. Bodkin et al. (2004) examined dive distributions of 14 foraging northern sea otters in Port Althorp, southeast Alaska, from May through July. This study also found that dive behavior was related to sex and size, with females averaging a maximum dive depth of 49 m compared to 82 m for larger males (Bodkin et al. 2004). Wolt (2014) also examined the dive behavior of sea otters in Alaska, but focused primarily on those with dependent pups.

Thometz et al. (2016) found that the mean foraging dive depth for all otters was 8.3 m and the range of mean dives was 2.3 to 28.6 m. Bodkin et al. (2004) calculated that sea otters in their southeast Alaska study completed a mean of 174 foraging dives per day and that these dives averaged a depth of 18.9 m (SD = 4.6 m) (Bodkin et al. 2004). Foraging dives had a mean depth of 2.7 m (SD = 0.2 m) when compared to traveling dives by the same animals (Bodkin et al. 2004). An observational study of sea otters off northern Washington state (Laidre and Jameson 2006) reported an average foraging dive time of 0.9 min (54 seconds), consistent with both Bodkin et al. (2004) and Thometz et al. (2016). Finally, Wolt et al. (2012) looked at sea otter dive behavior in Simpson Bay, Alaska where dives mirrored the bathymetry of the bay, which was roughly 30 m deep. Mean dive duration, depending on bathymetry, ranged from 62 to 114 seconds for individual foraging dives (Bodkin et al. 2004; Thometz et al. 2016; Wolt et al. 2012).

While dive data is available for foraging sea otters in southeast Alaska and coastal California, these foraging dives have been found to mirror the bathymetry of the local area. According to all existing dive data, sea otters may be found in up to 100 m of water (Bodkin et al. 2004; Thometz et al. 2016),

depending on bathymetry in the areas where dive data was logged. In the NWTT Study Area, off the coast of Washington, sea otters would occupy similar depths. While most sea otters would forage in waters less than 40 m deep (Bodkin et al. 2004; Thometz et al. 2016; Wolt et al. 2012), it is assumed any sea otter found in more than 40 m of water would be diving to or near the bottom in order to forage, unless they are traveling, grooming, or participating in social interactions.

While several studies have created activity budgets for sea otters (Bodkin et al. 2007; Esslinger et al. 2014; Wolt 2014; Yeates et al. 2007), the data from Laidre et al. (2009) is the most robust, utilizing the behavior of 25 sea otters, and was taken from observations in coastal Washington, within the NWTT Study Area. According to Laidre et al. (2009), within a 24-hour time budget where all age and sex classes of sea otters were combined, sea otters spent roughly 45 percent of their time resting, 41 percent of their time foraging, and 14 percent in “other” behavior. Laidre et al. (2009) classified “other” behavior (similar to (Gelatt et al. 2002)), as grooming, swimming, or any active non-feeding behavior. As discussed above, Bodkin et al. (2004) describe this type of behavior as “traveling dives” and measured these as averaging around 2.7 m.

Sea otters are expected to forage at or near the seafloor; therefore, their dive depth is dictated by the bathymetry of the location (Wolt et al. 2012) in their typical coastal habitats. In order to build a representative depth distribution for the northern sea otter, the activity budget in Laidre et al. (2009) was used and adjusted for time spent underwater. The 0-5 m bin includes time spent underwater while traveling, participating in “other” behavior, and foraging between 0-5 m. The total percentage of time spent in the 0-5 m depth bin was calculated to be 29.2 percent of underwater time. The remaining 70.8 percent of underwater time would be expected to be at depths between 6 and 100 m (accounting for depths greater than the deepest recorded otter dive), and dependent on the water depth of the foraging area. The depth distribution of northern sea otters is given in Table 19.

Table 19. Percentage of Time at Depth for the Northern Sea Otter

Depth Bin (m)	% of Time in Other (Traveling) Behavior at Depth	% of Time Foraging at Depth	% of Time at Depth
0-5	25.5	3.7	29.2
6-100	0	70.8	70.8

2.2.2.2 Family Otariidae

2.2.2.2.1 *Arctocephalus townsendi*, Guadalupe Fur Seal

The Guadalupe fur seal breeds at only two rookery locations: the east side of Isla Guadalupe and the southeast side of Isla Bonito del Este (Maravilla-Chavez and Lowry 1999). Single animals have occasionally been seen at islands along the coast of Baja, California and California (Stewart et al. 1987). Squid, teleost fish, and crustaceans comprise the diet of the Guadalupe fur seal (Aurióles-Gamboá and Camacho-Ríos 2007; Belcher and Lee 2002; Riedman 1990).

Little data have been collected on the dive behavior of Guadalupe fur seals. In order to build a representative depth distribution for Guadalupe fur seals, data from Figure 2 in Lander et al. (2000) were used. Lander et al. (2000) obtained information on the diving behavior of Guadalupe fur seals by satellite tagging a single female following rehabilitation off southern California. Most dives were performed at night and lasted 2–4 min (mean dive time of 1.7 min, SD=1.0 min), with a maximum dive duration of 18 min (Lander et al. 2000). The mean dive depth was 15.7 m (SD=11.8 m). The maximum

recorded dive depth was 130 m, although most dives were less than 20 m deep. While Lander et al. (2000) reported only the number of dives to various depth bins, this is used as a proxy for the time spent at depth. The bins over 40 m were presented together as they represented a small percentage of time at depth. The depth distribution for Guadalupe fur seals is given in Table 20.

Table 20. Percentage of Time at Depth for the Guadalupe Fur Seal¹

Depth Bin (m)	% of Time at Depth
0–20	92.0
20–40	7.0
40–160	1.0

¹Based on data from Lander et al. (2000)

Data from Gallo-Reynoso et al. (2008) also studied the diving behavior of one adult female, finding a mean dive depth of 17 m (SD=10 m) with an average dive duration of 2.6 min (SD=1.4 min).

2.2.2.2 *Callorhinus ursinus*, Northern Fur Seal

Northern fur seals occupy the pelagic waters of the North Pacific Ocean, Bering Sea, and Sea of Japan, ranging coastally as far south as Baja, California, Mexico, and Japan, and with an at-sea southern limit around 35°N (Gelatt and Lowry 2008). Northern fur seals are known to feed in the deep waters along the continental shelf break, as well as in shallower waters of the shelf itself (Gentry 2002; Ponganis et al. 1992). The diet of northern fur seals varies regionally and seasonally, but it is comprised principally of finfish (e.g., Pacific herring, sand lance, capelin, myctophids) and squid; they will occasionally feed upon other prey, such as birds and crustaceans (Ream et al. 2005; Riedman and Estes 1990).

In order to build a representative depth distribution for the Northern fur seals, data from Table 1 in Kooyman et al. (1976) were used. The dive behavior and physiology of the northern fur seal was among the earliest pinniped species to be tagged and tracked (Kooyman et al. 1976). Table 1 in Kooyman et al. (1976) indicates the number of dives to specific depth bins; these data will be used as a proxy for percentage of time spent at depth. Kooyman et al. (1976) found that shallow dives (0–20 m) lasted less than 1 min in duration. Deeper dives, from 20–140 m, lasted from 2–5 min. The maximum dive depth in this study was 190 m. The average interval between dives was 17 min. The depth distribution for Northern fur seals is given in Table 21.

Table 21. Percentage of Time at Depth for the Northern Fur Seal¹

Depth Bin (m)	% of Time at Depth
0–20	48.36
20–50	42.32
50–80	5.94
80–110	0.96
110–140	1.99
140–170	0.39
170–200	0.04

¹Based on data from Kooyman et al. (1976)

While other studies did not include depth distributions for northern fur seals, they did provide additional information to categorize dive behavior. Because northern fur seals spend the vast majority of their lives at sea (87–90 percent of the year), only coming ashore to breed for 35–45 days from June-to-August, most of the tagging studies examine the dive behavior of females on excursions from

breeding colonies in the Bering Sea (Baker 2007; Baker and Donohue 2000; Gentry 2002; Goebel et al. 1991; Kooyman et al. 1976; Kuhn et al. 2009; Kuhn et al. 2010; Ream et al. 2005). In general, adult females on foraging excursions generally follow one of three dive profiles: shallow, deep, or mixed depth. Shallow-diving seals show a crepuscular pattern with dive depths varying according to movement of the deep scattering layer. Deep-diving seals show no temporal pattern, apparently ignoring the diel movements of vertically migrating prey, and have no consistent change in depth within bouts. Mixed-depth divers alternate between dive profiles, perhaps shifting prey types (Gentry et al. 1986). Female northern fur seals dive mostly at night (68 percent) (Gentry et al. 1986). Individuals may be consistent in their diving behavior, presumably choosing prey sources at different depths, as evidenced by unique fatty acid profiles specific to a differentiated prey type (deep versus vertically migrating species, for example); however, other evidence points towards a seasonal shift in dive behavior (Gentry 2002; Gentry et al. 1986; Hobson and Sease 1998; Hobson et al. 1997). Radio-tracking studies suggest that deep-diving patterns are used while foraging on the continental shelf, while shallow-diving patterns occur over deeper waters off the shelf break or in the Aleutian Basin (Gentry et al. 1986; Goebel et al. 1991).

Female northern fur seals have been recorded diving to a maximum depth of 256 m (Ponganis et al. 1992), although they most frequently dive between 50 and 60 m (Gentry et al. 1986). The activity budgets for adult females on foraging trips was 17 percent of time spent resting, 26 percent of time spent diving, and 57 percent of time spent in surface active time (Gentry et al. 1986).

Male northern fur seal diving behavior has also been examined (Sterling and Ream 2004) and has revealed that like parturient females, juvenile males exhibit shallow versus deep dive patterns based upon foraging location (deep dives in water less than 200 m deep, shallow nighttime dives in waters up to 3,000 m deep). In a study of 19 juvenile male northern fur seals on foraging excursions during the breeding season in the Bering Sea, the maximum recorded dive depth was 175 m, with a mean dive depth of 17.5 m (SD=1.5 m) (Sterling and Ream 2004).

2.2.2.2.3 *Eumetopias jubatus*, Steller Sea Lion

Steller sea lions inhabit the Pacific Ocean north of approximately 30°N latitude (Loughlin 2002; Schusterman 1981). They feed on an assortment of shallow water fish, cephalopods, bivalves, and crustaceans (Schusterman 1981; Tollit et al. 2004).

Rehberg et al. (2009) reported on the summer diving behavior of five adult sea lions belonging to the Eastern Alaska (E AK) stock. They also summarized similar data on four Western Alaska (W AK) stock females from Merrick (1995) and Merrick and Loughlin (1997). A dive was counted as any movement of the animal below 4 m depth. Visual inspection of Figure 2 and the text from Rehberg et al. (2009) results in the distribution of dive depths in Table 22.

Table 22. Dive Depth Distribution Data from Rehberg et al. (2009), Merrick (1995), and Merrick and Loughlin (1997) Used to Estimate Depth Distribution for the Steller Sea Lion

Depth Bin (m)	Percentage of Dives		
	E AK Stock	W AK Stock	Average
4-10	45	35	40
10-20	14	46	30
20-50	23.5	15	19.25
50-100	14	3	8.5
100-250	3.5	1	2.25
>250	0	0	0

Due to the lack of data on time spent at depth, the data in Table 22 will be used as a proxy for depth distribution data. Skinner et al. (2012) and Rehberg et al. (2009) reported that on average females spent 15.1 percent and 22.1 percent of their time at sea submerged, respectively. Therefore, the distribution in Table 22 will be adjusted to represent 22.1 percent of an animal's time, with 77.9 percent of time spent between depths of 0-4 m. The maximum dive depth reported in this study was 236 m, and the last dive bin is adjusted to reflect this. The resulting depth distribution is given in Table 23.

Table 23. Percentage of Time at Depth for the Stellar Sea Lion¹

Depth Bin (m)	% of Time at Depth
0-4	77.9
4-10	8.8
10-20	6.6
20-50	4.3
50-100	1.9
100-236	0.5

¹Based on data from Rehberg et al. (2009), Merrick (1995), and Merrick and Loughlin (1997)

The depth distribution in Table 23 is consistent with research suggesting most dives are within the top 50 m (Fadely et al. 2005; Loughlin et al. 1998; Loughlin et al. 2003; Merrick and Loughlin 1997; Merrick et al. 1994; Skinner et al. 2012). It is shallower than the maximum reported dive depth of 328 m by a juvenile sea lion (Loughlin et al. 2003). There is evidence that adult females dive deeper during the winter than the summer (Merrick and Loughlin 1997; Merrick et al. 1994), although in both seasons the majority of dives are shallow (<50 m).

2.2.2.2.4 *Zalophus californianus*, California Sea Lion

California sea lions primarily breed on island beaches off southern California; along Baja California, Mexico, and in the Gulf of California (Heath 2002). They eat a variety of prey, including schooling fish, crustaceans, and cephalopods (García-Rodríguez and Aurióles-Gamboa 2004; Melin et al. 2008; Porrás-Peters et al. 2008). California sea lions often make shallow dives in some coastal areas for epipelagic prey and deeper dives in others, such as in the Gulf of California for mesopelagic prey (Costa et al. 2004; Lowry and Carretta 1999; Melin and DeLong 1999).

In order to build a representative depth distribution for California sea lions, data from Figure 4 in Feldkamp et al. (1989) and Figure 1a in Weise (2006) were used. Both figures show the number of dives to specific depth bins; thus, the data from Feldkamp et al. (1989) and Weise (2006) will be used as a

proxy for percentage of time spent at depth. Because data indicate differences in diving behavior between males (Weise 2006) and females (Feldkamp et al. 1989; Kuhn 2006; Melin et al. 2008), data from Weise (2006) and Feldkamp et al. (1989) were averaged to create a composite distribution. Weise (2006) reported an average dive depth of 32 m, but most dives occurred within the depth range of 10 to 20 m. Feldkamp et al. (1989) found that the maximum dive depth for females was roughly 274 m. The mean depths ranged from 31-to-98 m, but the majority of dives went to between 20 and 50 m. In order to account for the time spent at the surface, an activity budget from Table 3 in Thomas et al. (2010) was used, incorporating both male and female sea lions. Based on these numbers, a percent time swimming at the surface (63.62 percent) was added into the average taken from the Feldkamp et al. (1989) and Weise (2006) data. The remaining averages from both studies were redistributed proportionally to account for the remaining 36.38 percent of time. The resulting depth distribution profile includes data from all three studies. The depth distribution for California sea lions is given in Table 24.

Table 24. Percentage of Time at Depth for the California Sea Lion¹

Depth Bin (m)	% of Time at Depth	Depth Bin (m)	% of Time at Depth
0–10	68.3	120–130	0.5
10–20	5.9	130–140	0.4
20–30	6	140–150	0.5
30–40	4.8	150–160	0.5
40–50	4	160–170	0.5
50–60	2.3	170–180	0.3
60–70	1.5	180–190	0.1
70–80	1	190–200	0.1
80–90	0.7	200–210	0.2
90–100	0.7	210–220	0.1
100–110	0.7	220–230	0.1
110–120	0.6	230–500	0.2

¹Based on data from Feldkamp et al. (1989), Weise (2006), and Thomas (2010)

While other studies did not include usable depth distributions for California sea lions, they did provide additional information to categorize dive behavior. Mean dive depths reported for female California sea lions range from 45–70 m with maximum dive depths ranging from 104–279 m (Kuhn 2004; Kuhn 2006; Melin et al. 2008). While these dataset does not necessarily capture the maximum dive depths reported in several locations (greater than 200 m) (Costa et al. 2004; Feldkamp et al. 1989; Kuhn 2004; Kuhn 2006; Melin et al. 2008), the depth distribution is consistent with studies by Kuhn (2004) and Melin (2008), both of whom demonstrated that the majority of dives (ranging from 55–85 percent) were less than 50 m deep. Similarly, Melin and DeLong (1999) reported that most dives were shallower than 100 m; therefore, although the depth distribution in Table 24 may be shallower than seen in some locations, it does capture published diving behavior for the species and is representative for California sea lions.

2.2.2.3 Family Phocidae

2.2.2.3.1 *Mirounga angustirostris*, Northern Elephant Seal

Northern elephant seals are limited to the North Pacific Ocean, with breeding haul outs located along the western coast of North America from northern California to Baja California and Mexico. Seals utilize deep waters for foraging, traveling north and west from breeding beaches and traveling as far north as the Gulf of Alaska and Aleutian Islands (Hindell 2002). Elephant seals feed primarily on vertically

migrating epipelagic and mesopelagic squid, but they eat a variety of prey species, including elasmobranchs, crustaceans, cephalopods, and fish (DeLong and Stewart 1991; Sinclair 1994; Stewart and Huber 1993).

In order to build a representative depth distribution for Northern elephant seals, data from DeLong and Stewart (1991), Hakoyama et al. (1994), Le Boeuf et al. (1986), and Naito et al. (1989) were used. In addition, a surface time from DeLong and Stewart (1991) was incorporated into these data. The text of DeLong and Stewart (1991) states that Northern elephant seals are submerged 86 percent of the time, leaving an above-surface time percentage of 14 percent (DeLong and Stewart 1991). Due to the lack of data on time spent at depth, the data from these sources, citing percentage of dives to specific depth bins will be used as a proxy for percentage of time spent at depth. As these data show sex differences in foraging and diving behavior (DeLong and Stewart 1991; Hakoyama et al. 1994; Le Boeuf et al. 1986; Le Boeuf et al. 1993; Naito et al. 1989), males and females were averaged together to create a representative depth distribution. In males, the data averaged from Figure 1 in DeLong and Stewart (1991) and Figure 4 in Hakoyama et al. (1994) showed that 83 percent of dives took place between the surface and 500 m, most frequently occurring from 300–500 m. In females, these data averaged from Figure 1 in Le Boeuf et al. (1986), Figure 2 in Naito et al. (1989), and Figure 4 in Hakoyama et al. (1994) showed that 85 percent of dives were to depths of 300–700 m, most frequently from 400–600 m. For males, dives deeper than 500 m were unusual, whereas dives over 700 m were rare for females (DeLong and Stewart 1991; Hakoyama et al. 1994; Le Boeuf et al. 1988; Le Boeuf et al. 1986; Naito et al. 1989). Data from each figure were summed into 100 m bins and averaged together. The remaining bins of each study were averaged and redistributed proportionally to account for the remaining 86 percent of non-surface time. The resulting depth distribution profile includes data from all studies. The depth distribution for Northern elephant seals is given in Table 25.

Table 25. Percentage of Time at Depth for the Northern Elephant Seal¹

Depth Bin (m)	% of Time at Depth
0–100	21.4
100–200	7.6
200–300	8.9
300–400	13.4
400–500	18
500–600	16.3
600–700	10
700–800	2.2
800–900	0.8
900–1000	0.5
1,000–1,100	0.5
1,100–1,200	0.4

¹Based on data from DeLong and Stewart (1991), Hakoyama et al. (1994), Le Boeuf et al. (1986), and Naito et al. (1989)

While other studies did not include usable depth distributions for northern elephant seals, they did provide additional information to categorize dive behavior. Robinson et al. (2012) found that the dive depths of most seals showed a clear diel pattern, consistent with targeting vertically migrating prey species. Robinson et al. (2012) found the mean dive depth to be 516 m and a mean dive duration of 23.1 min. Active-bottom dives made up the greatest percentage of dives (54.0 percent) in this study (Robinson et al. 2012). Maximum dive depths for elephant seals covered a wide range of depths, from

under 1,000 m (Crocker et al. 2006; Davis et al. 2001; Davis and Weihs 2007; Le Boeuf et al. 1988) to the deepest dive ever recorded by an elephant seal at 1,747 m (Robinson et al. 2012).

2.2.2.3.2 *Phoca vitulina*, Harbor Seal

Harbor seals are found in shallow inshore and coastal waters of the Northern Hemisphere (Burns 2002). Prey species include epibenthic and benthic fish (e.g., sandlance, flounder, and herring) and squid (Brown and Mate 1983; Olesiuk 1993; Payne and Selzer 1989).

In order to build a representative depth distribution for harbor seals, data from Table 3 in Womble et al. (2014) and an activity budget from Bowen et al. (1999) were used. Womble et al. (2014) studied the dive behavior of 12 female harbor seals in Alaska. Table 3 shows the percentage of dives to specific depth bins; thus, these data from Womble et al. (2014) will be used as a proxy for percentage of time spent at depth. Seals in this study were found to dive most frequently (81.6 percent) to depths shallower than 50 m. It should be noted that this study cites the percentage of dives to certain depths, as opposed to the percentage of time spent in each depth bin, including time spent in that bin on the way to a deeper bin. However, the Womble et al. (2014) data are the best available estimate of time spent at depth. The activity budget from Bowen et al. (1999) provided a surface time of 85.01 percent. The remaining bins from Womble et al. (2014) were redistributed proportionally to account for the remaining 14.99 percent of non-surface time. The resulting depth distribution profile includes data from both studies. The depth distribution for harbor seals is given in Table 26.

Table 26. Percentage of Time at Depth for the Harbor Seal¹

Depth Bin (m)	% of Time at Depth
0-4	85.01
4-50	12.22
50-100	2.32
100-150	0.37
150-200	0.05
200-250	0.02
250-300	0.01

¹Based on data from Womble et al. (2014) and Bowen et al. (1999)

While other studies did not include usable depth distributions for harbor seals, they did provide additional information to categorize dive behavior. Gjertz et al. (2001) reported the maximum depth reached by harbor seals to be within the 200–350 m range; however, additional studies have reported shallower maximum dive depths, reaching less than 100 m off of Nova Scotia (Bowen et al. 1999), Svalbard (Jorgensen et al. 2001), and in Prince William sound (Frost et al. 2001). Hastings et al. (2004) observed that less than six percent of harbor seals dive to depths greater than 100 m, although one seal dove to 508 m (Hastings et al. 2004). Harbor seals near Svalbard, Norway, dove to a maximum depth of 452 m, although 50 percent of dives were shallower than 40 m, and 95 percent were to less than 250 m (Gjertz et al. 2001). Eguchi and Harvey (2005) observed that males dive deeper than females, with males diving to depths shallower than 154 m and females diving to depths shallower than 76 m, 95 percent of the time.

2.2.3 Sea Turtles

2.2.3.1 Family Dermochelyidae

2.2.3.1.1 *Dermochelys coriacea*, Leatherback Sea Turtle

The leatherback turtle is globally distributed in tropical, subtropical, and warm-temperate waters throughout the year, and throughout cooler temperate waters during warmer months (James et al. 2005a; National Marine Fisheries Service (NMFS) and United States Fish and Wildlife Service (USFWS) 1993). Adult leatherback turtles forage in temperate and subpolar regions in all oceans and migrate to tropical nesting beaches between 30° N and 20° South (Bleakney 1965; Brongersma 1972; Goff and Lien 1988; Threlfall 1978). The leatherback is typically associated with continental shelf habitats and pelagic environments. To a large extent, the oceanic distribution of leatherbacks may reflect the distribution and abundance of their favored prey; macro-planktonic soft-bodied animals such as jellyfish, salps, and pyrosomes (Wallace et al. 2013). It is suggested that leatherbacks make scouting dives while transiting as an efficient means for sampling prey density and perhaps also to feed opportunistically at these times (James et al. 2006b; Jonsen et al. 2007).

In order to build a representative depth distribution for leatherback turtles, data from Figure 2 as well as the text of Houghton et al. (2008) were used. Houghton et al. (2008) tagged 13 adult leatherback sea turtles at two sites over the course of four years. The data in Figure 2 accounts only for the percentage of dives made to each depth bin, rather than the percent of time spent in each bin. The text states the percent of dives were made to less than 10 m (18.9 percent), which is considered the surface bin. This representative depth distribution was created to account for several behavioral states, including post-nesting, migration, and foraging. The mean dives in this study ranged from 32.5–69.0 m. Houghton et al. (2008) found that 99.6 percent of dives were to depths shallower than 300 m with only 0.4 percent extending to greater depths. These findings support the hypothesis that deep dives are periodically employed to survey the water column for diurnally descending gelatinous prey. The depth distribution for leatherback turtles is given in Table 27.

Table 27. Percentage of Time at Depth for the Leatherback Turtle¹

Depth Bin (m)	% of Time at Depth
0–10	18.93
10–100	65.26
100–200	14.63
200–300	0.818
300–400	0.119
400–500	0.103
500–600	0.069
600–700	0.023
700–800	0.015
800–900	0.015
900–1,000	0.007
1,000–1,100	0.004
1,100–1,200	0.001
1,200–1,280	0.006

¹Based on data from Houghton et al. (2008)

While other studies did not include usable depth distributions for leatherback turtles, they did provide additional information to categorize dive behavior. The leatherback is the deepest diving sea turtle, with a recorded maximum depth of 1,280 m (Doyle et al. 2008), though most dives are much shallower, usually less than 250 m (Hays et al. 2004; Sale et al. 2006). Fossette et al. (2007) reported that eighty percent of the leatherback's time at sea is spent diving, which is in agreement with the roughly 20 percent of time in the surface bin of the representative depth distribution. (Dodge et al. 2014) found that over 90 percent of the time was spent in the top 100 m of the water column, and 25 percent of time was spent at the surface. Similarly, in the Atlantic, Hays et al. (2004) determined that leatherbacks spent 71–94 percent of their diving time at depths from 70 to 110 m. Daytime foraging dives off of the Canadian Atlantic coast during summer ranged between 5.5 and 97 m with a median depth of 21.5 m (Heaslip et al. 2012). Leatherbacks dive deeper and longer in the lower latitudes versus the higher (James et al. 2005a; James et al. 2005b), where they are known to dive to waters with temperatures just above freezing (James et al. 2006a; Jonsen et al. 2007). James et al. (2006b) noted that dives in higher latitudes are punctuated by longer surface intervals and much more time at the surface; individuals spend up to 50 percent of their time at or near the surface in northern foraging areas, perhaps in part to thermoregulate (i.e., bask). While transiting, leatherbacks make longer and deeper dives (Jonsen et al. 2007). During inter-nesting periods, tag data has revealed that dives are likely constrained by bathymetry adjacent to nesting sites (Hays et al. 2006; Myers and Hays 2006).

2.3 Conclusions

The recommended static depth distributions are provided for 32 marine animal species or guilds occurring within the NWTT Study Area. These distributions, especially those that rely on surrogates, should be updated periodically as new data become available. Also, for most species, only a single depth distribution is presented; ideally, each species should have multiple distributions available, depending on the behavior and age/sex class of the animals being modeled, as well as the geographic location and season in which the simulation occurs. More detailed depth distribution data will permit improved realism for the scenarios being modeled.

This page left intentionally blank.

3. Marine Mammal Group Size Information

Many marine mammals are known to travel and feed in groups. NAEMO accounts for this behavior by incorporating species-specific group sizes into the modeled animal distributions, and accounting for statistical uncertainty around the group size estimate. Methods for determining group size varied between regions of the NWTT Study Area, based on data availability and the recommendations of the research groups that provided density information.

Group size data were determined for four regions of the NWTT Study Area: inland waters of Puget Sound, the San Juan Islands/Strait of Juan de Fuca, offshore waters, and the Behm Canal in southeastern Alaska. For species in the offshore waters portion of the NWTT Study Area, survey information was used. Group size values were calculated from raw sightings data from The National Marine Fisheries Service Southwest Fisheries Science Center survey cruises (1996-2004) (Barlow 2016; Barlow and Forney 2007; Barlow et al. 2010). For most species, only sightings north of 38°N were included in the analysis. However, due to low sighting numbers, sightings data from the entire California Current Ecosystem region were used for minke whales, gray whales, and *Kogia* spp.

For all other regions in the NWTT Study Area, group size data were collected through a comprehensive and systematic review of available visual survey data and relevant literature. Journal articles, books, technical reports, cruise reports, funding agency reports, theses, dissertations, and raw data from individual researchers, theses, and dissertations were assessed for this report.

Group size data include mean group size and SD. The SDs are incorporated in NAEMO by randomly and repetitively selecting a value from the Poisson or lognormal distribution defined by the mean group size and standard deviation provided. The National Marine Fisheries Service Southwest Fisheries Science Center provided recommendations on which distribution to use for each species. Group size data for the four regions of the NWTT Study Area are given in Tables 28 through 30. Pinnipeds and turtles are primarily observed individually while at sea, and therefore are assigned a group size of 1.

Table 28. Mean Group Size, Standard Deviation, and Ranges for Marine Species in the Offshore Waters Portion of the NWTT Study Area

Common and Scientific Name	Mean Group Size	SD	Distribution	References
Cetaceans				
Family Balaenopteridae				
Minke whale <i>Balaenoptera acutorostrata</i>	1.09	0.36	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Sei whale <i>Balaenoptera borealis</i>	2.08	1.29	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Bryde's whale <i>Balaenoptera edeni</i>	1	--	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Blue whale <i>Balaenoptera musculus</i>	1.67	1.16	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)

Table 28. Mean Group Size, Standard Deviation, and Ranges for Marine Species in the Offshore Waters Portion of the NWTT Study Area (Cont'd)

Common and Scientific Name	Mean Group Size	SD	Distribution	References
Cetaceans				
Family Balaenopteridae				
Fin whale <i>Balaenoptera physalus</i>	1.93	1.6	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Humpback whale <i>Megaptera novaeangliae</i>	1.88	1.48	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Family Delphinidae				
Short-beaked common dolphin <i>Delphinus delphis</i>	175.88	258.87	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Short-finned pilot whale <i>Globicephala macrorhynchus</i>	15.93	5.62	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Risso's dolphin <i>Grampus griseus</i>	23.72	31.33	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Pacific white-sided dolphin <i>Lagenorhynchus obliquidens</i>	64.79	164.16	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Northern right whale dolphin <i>Lissodelphis borealis</i>	36.25	63.89	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Killer whale <i>Orcinus orca</i>	6.95	5.11	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Striped dolphin <i>Stenalla coeruleoalba</i>	36	58.65	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Bottlenose dolphin <i>Tursiops truncatus</i>	9.37	8.04	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Family Eschrichtiidae				
Gray whale <i>Eschrichtius robustus</i>	2.07	1.85	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Family Kogiidae				
<i>Kogia</i> spp.	1.97	2.03	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)

Table 28. Mean Group Size, Standard Deviation, and Ranges for Marine Species in the Offshore Waters Portion of the NWTT Study Area (Cont'd)

Common and Scientific Name	Mean Group Size	SD	Distribution	References
Cetaceans				
Family Phocoenidae				
Dall's porpoise <i>Phocoenoides dalli</i>	3.67	3.02	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Harbor porpoise <i>Phocoena phocoena</i>	2.42	4.97	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Family Physeteridae				
Sperm whale <i>Physeter macrocephalus</i>	5.95	7.4	Log-normal	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Family Ziphiidae				
Baird's beaked whale <i>Berardius bairdii</i>	7.72	6.7	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)
Small beaked whale guild (<i>Ziphius</i> and <i>Mesoplodon</i> spp.)	2.2	1.19	Poisson	Barlow (2016), Barlow et al. (2010), Barlow and Forney (2007)

Table 29. Mean Group Size, Standard Deviation, and Range for Marine Species in the Inland Waters of Puget Sound and San Juan Islands/Strait of Juan de Fuca Portion of the NWTT Study Area

Common and Scientific Name	San Juan Islands/Strait of Juan de Fuca		Puget Sound		Distribution	References
	Mean Group Size	SD	Mean Group Size	SD		
Cetaceans						
Family Balaenopteridae						
Minke whale <i>Balaenoptera acutorostrata</i>	1.59	0.54	1.21	0.1	Poisson	Orca Network (2017), Cascadia Research Collective (2017), Smultea (2017)
Humpback whale <i>Megaptera novaeangliae</i>	2.17	2.19	1.33	0.14	Poisson	Orca Network (2017), Cascadia Research Collective (2017)
Family Delphinidae						
Pacific white-sided dolphin <i>Lagenorhynchus obliquidens</i>	28.5	13.44	N/A	N/A	Log-normal	Orca Network (2017), Cascadia Research Collective (2017)
Killer whale, Southern Resident <i>Orcinus orca</i>	11.38	4.78	7.12	2.68	Log-normal	Orca Network (2017), Cascadia Research Collective (2017), Smultea (2017), The Whale Museum (2017b)
Killer whale, West Coast Transient <i>Orcinus orca</i>	7.12	3.5	5.8	1.12	Log-normal	Orca Network (2017), Cascadia Research Collective (2017), Houghton (2015), The Whale Museum (2017a)
Family Eschrichtiidae						
Gray whale <i>Eschrichtius robustus</i>	1.37	1.03	1.52	0.04	Poisson	Orca Network (2017), Cascadia Research Collective (2017)
Family Phocoenidae						
Harbor porpoise <i>Phocoena phocoena</i>	10.12	14.99	2.66	1.79	Poisson	Orca Network (2017), Cascadia Research Collective (2017), Jefferson (2016), Smultea (2017)
Dall's porpoise <i>Phocoenoides dalli</i>	3.52	3.55	5.04	0.76	Poisson	Orca Network (2017), Cascadia Research Collective (2017), Smultea (2017)

Table 30. Mean Group Size, Standard Deviation, and Range for Marine Species in the Behm Canal Portion of the NWTT Study Area

Common and Scientific Name	Mean Group Size	SD	Distribution	References
Cetaceans				
Family Balaenopteridae				
Minke whale <i>Balaenoptera acutorostrata</i>	1	0	Poisson	Dahlheim et al. (2009), Straley et al. (2018)
Fin whale <i>Balaenoptera physalus</i>	2.1	0.14	Poisson	Dahlheim et al. (2009)
Humpback whale <i>Megaptera novaeangliae</i>	1.72	0.49	Poisson	Dahlheim et al. (2009)
Family Delphinidae				
Pacific white-sided dolphin <i>Lagenorhynchus obliquidens</i>	43.78	55.14	Log-normal	Dahlheim et al. (2009)
Killer whale, Alaska Resident <i>Orcinus orca</i>	28.65	12.39	Log-normal	Dahlheim et al. (2009)
Killer whale, Northern Resident <i>Orcinus orca</i>	28.65	12.39	Log-normal	Dahlheim et al. (2009)
Killer whale, Offshore <i>Orcinus orca</i>	30.75	17.5	Log-normal	Dahlheim et al. (2009)
Killer whale, West Coast Transient <i>Orcinus orca</i>	5.88	3.64	Log-normal	Dahlheim et al. (2009)
Family Phocoenidae				
Harbor porpoise <i>Phocoena phocoena</i>	1.96	0.41	Poisson	Dahlheim et al. (2009)
Dall's porpoise <i>Phocoenoides dalli</i>	3.45	0.64	Poisson	Dahlheim et al. (2009)

This page left intentionally blank.

4. REFERENCES

- Acevedo-Gutiérrez, A., Croll, D. A., & Tershy, B. R. (2002). High Feeding Costs Limit Dive Time in the Largest Whales. *The Journal of Experimental Biology*, 205, 1747-1753.
- Aguilar, A. (2002). Fin Whale (*Balaenoptera physalus*). In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 435-438). San Diego, CA: Academic Press.
- Aguilar de Soto, N., Johnson, M., Madsen, P. T., Tyack, P. L., Bocconcelli, A., & Fabrizio Borsani, J. (2006). DOES INTENSE SHIP NOISE DISRUPT FORAGING IN DEEP-DIVING CUVIER'S BEAKED WHALES (ZIPHIUS CAVIROSTRIS)? *Marine Mammal Science*, 22(3), 690-699.
- Alves, F., Dinis, A., & Cascão, I. (2010). Bryde's Whale (*Balaenoptera brydei*) Stable Associations and Dive Profiles: New Insights into Foraging Behavior. *Marine Mammal Science*, 26(1), 202-212.
- Amano, M., & Yoshioka, M. (2003). Sperm Whale Diving Behavior Monitored Using a Suction-Cup-Attached TDR Tag. *Marine Ecology Progress Series*, 258, 291-295.
- Aoki, K., Amano, M., Mori, K., Kourogi, A., Kubodera, T., & Miyazaki, N. (2012). Active hunting by deep-diving sperm whales: 3D dive profiles and maneuvers during bursts of speed. *Marine Ecology Progress Series*, 444, 289-301.
- Aoki, K., Amano, M., Yoshioka, M., Mori, K., Tokuda, D., & Miyazaki, N. (2007). Diel Diving Behavior of Sperm Whales off Japan. *Marine Ecology Progress Series*, 349, 277-287.
- Archer II, F. I. (2002). Striped Dolphin: *Stenella coeruleoalba*. In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1201-1203). San Diego, CA: Academic Press.
- Arranz, P., de Soto, N. A., Madsen, P. T., Brito, A., Bordes, F., & Johnson, M. P. (2011). Following a foraging fish-finder: diel habitat use of Blainville's beaked whales revealed by echolocation. *PLoS one*, 6(12), e28353.
- Aurioles-Gamboa, D., & Camacho-Ríos, F. J. (2007). Diet and Feeding Overlap of Two Otariids, *Zalophus californianus* and *Arctocephalus townsendi*: Implications to Survive Environmental Uncertainty. *Aquatic Mammals*, 33(3), 315-326.
- Azzellino, A., Gaspari, S., Airoidi, S., & Nani, B. (2008). Habitat Use and Preferences of Cetaceans along the Continental Slope and the Adjacent Pelagic Waters in the Western Ligurian Sea. *Deep-Sea Research I*, 55, 296-323.
- Baird, R. W. (1994). *Foraging Behaviour and Ecology of Transient Killer Whales (Orcinus Orca)*. Doctor of Philosophy, Simon Fraser University.
- Baird, R. W., & Hanson, M. B. (1998). A Preliminary Analysis of the Diving Behavior of Dall's Porpoise in the Transboundary Waters of British Columbia and Washington. In Hill, P. S., Jones, B. & DeMaster, D. P. (Eds.), *Marine Mammal Protection Act and Endangered Species Act Implementation Program 1997* (pp. 99-110). Silver Spring, MD: National Marine Fisheries Service.
- Baird, R. W., Hanson, M. B., & Dill, L. M. (2005a). Factors Influencing the Diving Behaviour of Fish-eating Killer Whales: Sex Differences and Diel and Interannual Variation in Diving Rates. *Canadian Journal of Zoology*, 83, 257-267.

- Baird, R. W., Ligon, A. D., Hooker, S. K., & Gorgone, A. M. (2001). Subsurface and Nighttime Behaviour of Pantropical Spotted Dolphins in Hawai'i. *Canadian Journal of Zoology*, 79(6), 988-996.
- Baird, R. W., Martin, S. W., Webster, D. L., & Southall, B. L. (2014). *Assessment of Modeled Received Sound Pressure Levels and Movements of Satellite-Tagged Odontocetes Exposed to Mid-Frequency Active Sonar at the Pacific Missile Range Facility: February 2011 Through February 2013*. DTIC Document.
- Baird, R. W., McSweeney, D. J., Webster, D. L., Gorgone, A. M., & Ligon, A. D. (2003). *Studies of Odontocete Population Structure in Hawaiian Waters: Results of a Survey Through the Main Hawaiian Islands in May and June 2003*. (Contract Number AB133F-02-CN-0106). Seattle, WA: National Oceanic and Atmospheric Administration, Western Administrative Support Center. p. 24.
- Baird, R. W., Webster, D. L., McSweeney, D. J., Ligon, A. D., & Schorr, G. S. (2005b). *Diving Behavior and Ecology of Cuvier's (Ziphius cavirostris) and Blainville's Beaked Whales (Mesoplodon densirostris) in Hawai'i*. La Jolla, California: National Marine Fisheries Service (NMFS). p. 24.
- Baird, R. W., Webster, D. L., McSweeney, D. J., Ligon, A. D., Schorr, G. S., & Barlow, J. (2006). Diving Behaviour of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) Beaked Whales in Hawai'i. *Canadian Journal of Zoology*, 84, 1120-1128.
- Baird, R. W., Webster, D. L., Schorr, G. S., McSweeney, D. J., & Barlow, J. (2008). Diel Variation in Beaked Whale Diving Behavior. *Marine Mammal Science*, 24(3), 630-642.
- Baker, J. D. (2007). Post-weaning Migration of Northern Fur Seal *Callorhinus ursinus* Pups from the Pribilof Islands, Alaska. *Marine Ecology Progress Series*, 341, 243-255.
- Baker, J. D., & Donohue, M. J. (2000). Ontogeny of Swimming and Diving in Northern Fur Seal (*Callorhinus ursinus*) Pups. *Canadian Journal of Zoology*, 78, 100-109.
- Balcomb, K. C., III. (1989). Baird's Beaked Whale -- *Berardius bairdii* Stejneger, 1883: Arnoux's Beaked Whale -- *Berardius arnuxii* Duvernoy, 1851. In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals: Volume 4: River Dolphins and the Larger Toothed Whales* (Vol. 4, pp. 261-288). New York: Academic Press.
- Barlow, J. (2016). *Cetacean abundance in the California current estimated from ship-based line-transect surveys in 1991-2014*. La Jolla, CA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Southeast Fisheries Science Center (SEFSC), Marine Mammal and Turtle Division.
- Barlow, J., & Forney, K. A. (2007). Abundance and population density of cetaceans in the California Current ecosystem. *Fishery Bulletin*, 105(4), 509-526.
- Barlow, J., Henry, A., Redfern, J. V., Yack, T., Jackson, A., Hall, C., Archer, E., & Ballance, L. (2010). *Oregon, California and Washington line-transect and ecosystem (ORCAWHALE) 2008 cruise report*. La Jolla, CA: U.S. Department of Commerce, National Oceanic and Atmospheric Administration (NOAA), National Marine Fisheries Service (NMFS), Southwest Fisheries Science Center (SWFC), Protected Resources Division.
- Barlow, J., Tyack, P. L., Johnson, M. P., Baird, R. W., Schorr, G. S., Andrews, R. D., & de Soto, N. A. (2013). Trackline and point detection probabilities for acoustic surveys of Cuvier's and Blainville's beaked whales. *The Journal of the Acoustical Society of America*, 134(3), 2486-2496.

- Barrett-Lennard, L. G., Ford, J. K. B., & Heise, K. A. (1996). The Mixed Blessing of Echolocation: Differences in Sonar Use by Fish-Eating and Mammal-Eating Whales. *Animal Behaviour*, *51*, 553-565.
- Baumgartner, M., & Fratantoni, D. M. (2008). Diel periodicity in both sei whale vocalization rates and the vertical migration of their copepod prey observed from ocean gliders. *Limnol Oceanogr*, *53*, 2197-2209.
- Baumgartner, M., Lysiak, N. S. J., Schuman, C., Urban-Rich, J., & Wenzel, F. (2011). Diel vertical migration behavior of *Calanus finamarchicus* and its influence on right and sei whale occurrence. *Mar Ecol Prog Ser*, *423*, 167-184.
- Baumgartner, M. F. (1997). The Distribution of Risso's Dolphin (*Grampus griseus*) with Respect to the Physiography of the Northern Gulf of Mexico. *Marine Mammal Science*, *13*(4), 614-638.
- Beatson, E. (2007). The Diet of Pygmy Sperm Whales, *Kogia breviceps*, Stranded in New Zealand: Implications for Conservation. *Reviews in Fish Biology and Fisheries*, *17*, 295-303.
- Belcher, R. L., & Lee, T. E. J. (2002). *Arctocephalus townsendi*. *Mammalian Species*, *700*, 1-5.
- Best, P. B. (2001). Distribution and Population Separation of Bryde's Whale *Balaenoptera edeni* off Southern Africa. *Marine Ecology Progress Series*, *220*, 277-289.
- Bjorge, A., & Tolley, K. A. (2002). Harbor Porpoise. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 549-551). San Diego, CA: Academic Press.
- Black, N. A. (1994). *Behavior and Ecology of Pacific White-sided Dolphins (Lagenorhynchus obliquidens) in Monterey Bay, California*. (Master's thesis), San Francisco State University.
- Blanco, C., Aznar, J., & Raga, J. A. (1995). Cephalopods in the Diet of Striped Dolphin *Stenella coeruleoalba* from the Western Mediterranean during an Epizootic in 1990. *Journal of Zoology: Proceedings of the Zoological Society of London*, *237*, 151-158.
- Bleakney, J. S. (1965). Reports of Marine Turtles from New England and Eastern Canada. *Canadian Field-Naturalist*, *79*, 120-128.
- Blix, A. S., & Folkow, L. P. (1995). Daily Energy Expenditure in Free Living Minke Whales. *Acta Physiologica Scandinavica*, *153*(1), 61-66.
- Bloodworth, B. E., & Odell, D. K. (2008). *Kogia breviceps* (Cetacea: Kogiidae). *Mammalian Species*, *819*, 1-12.
- Bodkin, J. L., Esslinger, G. G., & Monson, D. H. (2004). Foraging Depths of Sea Otters and Implications to Coastal Marine Communities. *Marine Mammal Science*, *20*(2), 305-321.
- Bodkin, J. L., Monson, D. H., & Esslinger, G. G. (2007). Activity Budgets Derived From Time-Depth Recorders in a Diving Mammal. *The Journal of Wildlife Management*, *71*(6), 2034-2044.
- Bowen, W. D., Boness, D. J., & Iverson, S. J. (1999). Diving Behaviour of Lactating Harbour Seals and Their Pups During Maternal Foraging Trips. *Canadian Journal of Zoology*, *77*, 978-988.
- Brongersma, L. D. (1972). European Atlantic Turtles. In *Zoologische Verhandelingen* (Vol. 121, pp. 42-109). Leiden, Netherlands: E.J. Brill.
- Brown, R. F., & Mate, B. R. (1983). Abundance, Movements, and Feeding Habits of Harbor Seals, *Phoca vitulina*, at Netarts and Tillamook Bays, Oregon. *Fishery Bulletin*, *81*, 291-301.

- Brownell, R. L., Jr., Walker, W. A., & Forney, K. A. (1999). Pacific White-sided Dolphin -- *Lagenorhynchus obliquidens* Gill, 1865. In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 6: The Second Book of Dolphins and the Porpoises* (Vol. 6, pp. 57-84). New York: Academic Press.
- Budylenko, G. A. (1978). On Sei Whale Feeding in the Southern Ocean. *Reports of the International Whaling Commission*, 28, 379-385.
- Burns, J. J. (2002). Harbor Seal and Spotted Seal. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 552-560). San Diego, CA: Academic Press.
- Cacchione, D. A., Drake, D. E., Field, M. E., & Tate, G. B. (1987). Sea-Floor Gouges caused by Migrating Gray Whales off Northern California. *Continental Shelf Research*, 7(6), 553-560.
- Calambokidis, J., Schorr, G. S., Steiger, G. H., Francis, J., Bakhtiari, M., Marshall, G., Oleson, E. M., Gendron, D., & Robertson, K. (2008). Insights into the Underwater Diving, Feeding, and Calling Behavior of Blue Whales from a Suction-Cup-Attached Video-Imaging Tag (Cittercam). *Marine Technology Society Journal*, 41(4), 19-29.
- Cañadas, A., & Hammond, P. S. (2008). Abundance and Habitat Preferences of the Short-beaked Common Dolphin *Delphinus delphis* in the Southwestern Mediterranean: Implications for Conservation. *Endangered Species Research*, 4, 309-331.
- Cañadas, A., Sagarminaga, R., & García-Tiscar, S. (2002). Cetacean Distribution Related with Depth and Slope in the Mediterranean Waters off Southern Spain. *Deep-Sea Research I*, 49, 2053-2073.
- Cardinale, M., Casini, M., Arrhenius, F., & Håkansson, N. (2003). Diel spatial distribution and feeding activity of herring (*Clupea harengus*) and sprat (*Sprattus sprattus*) in the Baltic Sea. *Aquatic Living Resources*, 16, 283-292.
- Clapham, P. J. (2002). Humpback Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 589-592). San Diego, CA: Academic Press.
- Clarke, M. R. (2003). Production and control of sound by the small sperm whales, *Kogia breviceps* and *K. sima* and their implications for other Cetacea. *Journal of the Marine Biological Association of the UK*, 83, 241-263. doi: 10.1017/S0025315403007045h.
- Clarke, M. R., & Pascoe, P. L. (1985). The Stomach Contents of a Risso's Dolphin (*Grampus griseus*) Stranded at Thurlestone, South Devon. *Journal of Marine Biological Association of the United Kingdom*, 65, 663-665.
- Collective, C. R. (2017). *Orca Sightings Access Database 1980s-2015 provided by John Calambokidis on 7/28/2017* Olympia, WA: Cascadia Research Collective.
- Cooper, C. (1993). Diving Behaviour of Harbour Porpoises in the Lower Bay of Fundy. *Project Summary, Industry Services and Native Fisheries*, 42, 4.
- Costa, D. P., Kuhn, C., Weise, M., Shaffer, S., & Arnould, J. P. Y. (2004). When Does Physiology Limit the Foraging Behavior of Freely Diving Mammals. *International Congress Series*, 1275, 259-366.
- Crocker, D. E., Costa, D. P., Boeuf, B. J. L., Webb, P. M., & Houser, D. S. (2006). Impact of El Nino on the Foraging Behavior of Female Northern Elephant Seals. *Marine Ecology Progress Series*, 309, 1-10.

- Croll, D. A., Acevedo-Gutierrez, A., Tershy, B. R., & Urban-Ramirez, J. (2001). The Diving Behavior of Blue and Fin Whales: Is Dive Duration Shorter than Expected Based on Oxygen Stores? *Comparative Biochemistry and Physiology*, 129, 797-809.
- Cruickshank, R. A., & Brown, S. G. (1981). Recent Observations and Some Historical Records of Southern Right-Whale Dolphins *Lissodelphis peronii*. *Fisheries Bulletin*, 15, 109-121.
- Dahlheim, M. E., White, P. A., & Waite, J. M. (2009). Cetaceans of Southeast Alaska: Distribution and Seasonal Occurrence. *Journal of Biogeography*, 36, 410-426.
- Darling, J. D., Keogh, K. E., & Steeves, T. E. (1998). Gray Whale (*Eschrichtius robustus*) Habitat Utilization and Prey Species off Vancouver Island, B.C. *Marine Mammal Science*, 14(4), 692-720.
- Davis, R. W., & Fargion, G. S. (1996). *Distribution and Abundance of Cetaceans in the North-Central and Western Gulf of Mexico, Final Report. Volume II: Technical Report*. (OCS Study MMS 96-0027). New Orleans, LA: U.S. Dept of the Interior, Minerals Management Service. p. 356.
- Davis, R. W., Fargion, G. S., May, N., Leming, T. D., Baumgartner, M. F., Evans, W. E., Hansen, L. J., & Mullin, K. D. (1998). Physical Habitat of Cetaceans Along the Continental Slope in the North-Central and Western Gulf of Mexico. *Marine Mammal Science*, 14(3), 490-507.
- Davis, R. W., Fuiman, L. A., Williams, T. M., & Le Boeuf, B. J. (2001). Three-Dimensional Movements and Swimming Activity of a Northern Elephant Seal. *Comparative Biochemistry and Physiology Part A Molecular & Integrative Physiology*, 129A(4), 759-770.
- Davis, R. W., & Weihs, D. (2007). Locomotion in Diving Elephant Seals: Physical and Physiological Constraints. *Philosophical Transactions of the Royal Society of Britain*, 362, 2141-2150.
- De Vos, A., Clark, R. D., Johnson, C., Johnson, G., Kerr, I., Payne, R., & MADSEN, P. T. (2012). Cetacean sightings and acoustic detections in the offshore waters of Sri Lanka: March–June 2003. *J. Cetacean Res. Manage*, 12(2), 185-193.
- DeLong, R. L., & Stewart, B. S. (1991). Diving Patterns of Northern Elephant Seal Bulls. *Marine Mammal Science*, 7(4), 369-384.
- Dietz, R., Teilmann, J., Jorgensen, M.-P. H., & Jensen, M. V. (2002). *Satellite Tracking of Humpback Whales in West Greenland*. (NERI Technical Report No. 411). Roskilde, Denmark: National Environmental Research Institute. p. 40.
- DiMatteo, A., Jones, D., & Bartlett, B. (2017). *U.S. Navy Marine Species Density Database Phase III for the Atlantic Fleet Training and Testing Study Area*. Norfolk, VA: U. S. Department of the Navy (DON), U.S. Fleet Forces Command (USFF), Naval Facilities Engineering Command, Atlantic (NAVFACANT), Naval Undersea Warfare Center (NUWC).
- Dodge, K. L., Galuardi, B., Miller, T. J., & Lutcavage, M. E. (2014). Leatherback turtle movements, dive behavior, and habitat characteristics in ecoregions of the Northwest Atlantic Ocean. *PLoS One*, 9(3), e91726.
- Doksaeter, L., Olsen, E., Nottestad, L., & Ferno, A. (2008). Distribution and Feeding Ecology of Dolphins along the Mid-Atlantic Ridge between Iceland and the Azores. *Deep Sea Research II*, 55, 243-253.
- Dolphin, W. F. (1987a). Dive Behavior and Estimated Energy Expenditure of Foraging Humpback Whales in Southeast Alaska [USA]. *Canadian Journal of Zoology*, 65(2), 354-362.

- Dolphin, W. F. (1987b). Prey Densities and Foraging of Humpback Whales, *Megaptera novaeangliae*. *Experientia*, 43, 468-471.
- Domenici, P., Batty, R. S., Simila, T., & Ogam, E. (2000). Killer Whales (*Orcinus orca*) Feeding on Schooling Herring (*Clupea harengus*) Using Underwater Tail-Slaps: Kinematic Analyses of Field Observations. *The Journal of Experimental Biology*, 203, 283-294.
- Doniol-Valcroze, T., Lesage, V., Girard, J., & Michaud, R. (2011). Optimal foraging theory predicts diving and feeding strategies of the largest marine predator. *Behavioral Ecology*, 22(4), 880-888. doi: 10.1093/beheco/arr038.
- Dorsey, E. M., Stern, S. J., Hoelzel, A. R., & Jacobsen, J. (1990). Minke Whales (*Balaenoptera acutorostrata*) from the West Coast of North America: Individual Recognition and Small-scale Site Fidelity. *Reports of the International Whaling Commission*(Special Issue 12), 357-368.
- Doyle, T. K., Houghton, J. D. R., O'Súilleabháin, P. F., Hobson, V. J., Marnell, F., Davenport, J., & Hays, G. C. (2008). Leatherback Turtles Satellite-Tagged in European Waters. *Endangered Species Research*, 4, 23-31.
- Dunham, J. S., & Duffus, D. A. (2002). Diet of Gray Whales (*Eschrichtius robustus*) in Clayoquot Sound, British Columbia, Canada. *Marine Mammal Science*, 18(2), 419-437.
- Eguchi, T., & Harvey, J. T. (2005). Diving Behavior of the Pacific Harbor Seal (*Phoca vitulina richardii*) in Monterey Bay, California. *Marine Mammal Science*, 21(2), 283-295.
- Esslinger, G. G., Bodkin, J. L., Breton, A. R., Burns, J. M., & Monson, D. H. (2014). Temporal patterns in the foraging behavior of sea otters in Alaska. *Journal of Wildlife Management*, 689-700.
- Evans, W. E. (1974). Telemetry of Temperature and Depth Data From a Free Ranging Yearling California Gray Whale, *Eschrichtius robustus*. *Marine Fisheries Review*, 36(4), 52-58.
- Evans, W. E. (1975). *Distribution, Differentiation of Populations, and Other Aspects of the Natural History of Delphinus delphis Linnaeus in the Northeastern Pacific*. Doctor of Philosophy (Ph.D. Dissertation), University of California, Los Angeles, Los Angeles.
- Evans, W. E. (1994). Common Dolphin, White-Bellied Porpoise--*Delphinus delphis* Linnaeus, 1758. In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 5: The First Book of Dolphins* (Vol. 5, pp. 191-224). New York: Academic Press.
- Fadely, B. S., Robson, B. W., Sterling, J. T., Greig, A., & Call, K. A. (2005). Immature Steller Sea Lion (*Eumetopias jubatus*) Dive Activity In Relation to Habitat Features of the Eastern Aleutian Islands. *Fisheries Oceanography*, 14(1), 243-258.
- Feldkamp, S. D., DeLong, R. L., & Antonelis, G. A. (1989). Diving Patterns of California Sea Lions, *Zalophus californianus*. *Canadian Journal of Zoology*, 67, 872-883.
- Fernandez, R., Santos, M. B., Carrillo, M., Tejedor, M., & Pierce, G. J. (2009). Stomach Contents of Cetaceans Stranded in the Canary Islands 1996–2006. *Journal of the Marine Biological Association of the United Kingdom*, 89(5), 873-883.
- Fiscus, C. H., Rice, D. W., & Wolman, A. A. (1989). *Cephalopods from the Stomachs of Sperm Whales taken off California*. (NOAA Technical Report NMFS 83). National Oceanic and Atmospheric Administration, National Marine Fisheries Service. pp. 1-12.

- Fitch, J. E., & Brownell, R. L., Jr. (1968). Fish Otoliths in Cetacean Stomachs and Their Importance in Interpreting Feeding Habits. *Journal of the Fisheries Research Board of Canada*, 25(12), 2561-2574.
- Flinn, R. D., Trites, A. W., Gregr, E. J., & Perry, R. I. (2002). Diets of Fin, Sei, and Sperm Whales in British Columbia: An Analysis of Commercial Whaling Records, 1963-1967. *Marine Mammal Science*, 18(3), 663-679.
- Ford, J. K. B. (2002). Killer Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 669-676). San Diego, CA: Academic Press.
- Ford, J. K. B., Ellis, G. M., Barrett-Lennard, L. G., Morton, A. B., Palm, R. S., & III, K. C. B. (1998). Dietary Specialization in Two Sympatric Populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Canadian Journal of Zoology*, 76, 1456-1471.
- Forney, K. A., & Barlow, J. (1998). Seasonal Patterns in the Abundance and Distribution of California Cetaceans, 1991-1992. *Marine Mammal Science*, 14(3), 460-489.
- Fossette, S., Ferraroli, S., Tanaka, H., Ropert-Coudert, Y., Arai, N., & Sato, K. (2007). Dispersal and Dive Patterns in Gravid Leatherback Turtles during the Nesting Season in French Guiana. *Marine Ecology Progress Series*, 338, 233-247.
- Friedlaender, A. S., Hazen, E. L., Nowacek, D. P., Halpin, P. N., Ware, C., Weinrich, M. T., Hurst, T., & Wiley, D. (2009a). Diel Changes in Humpback Whale *Megaptera novaeangliae* Feeding Behavior in Response to Sand Lance *Ammodytes* spp. Behavior and Distribution. *Marine Ecology Progress Series*, 395, 91-100.
- Friedlaender, A. S., Lawson, G. L., & Halpin, P. N. (2009b). Evidence of Resource Partitioning Between Humpback and Minke Whales Around the Western Antarctic Peninsula. *Marine Mammal Science*, 25(2), 402-415.
- Frost, K. J., Simpkins, M. A., & Lowry, L. F. (2001). Diving Behavior of Subadult and Adult Harbor Seals in Prince William Sound, Alaska. *Marine Mammal Science*, 17(4), 813-834.
- Gallo-Reynoso, J. P., Figueroa-Carranza, A.-L., & Le Boeuf, B. (2008). Foraging behavior of lactating Guadalupe fur seal females. *Avances en el Estudio de los Mamíferos de México*, C. Lorenzo, E. Espinoza and J. Ortega (eds.). *Publicaciones Especiales*, 2, 595-614.
- García-Rodríguez, F. J., & Aurióles-Gamboa, D. (2004). Spatial and Temporal Variation In the Diet of the California Sea Lion (*Zalophus californianus*) in the Gulf of California, Mexico. *Fishery Bulletin*, 102(1), 47-62.
- Gelatt, T., & Lowry, L. (2008). *Callorhinus ursinus* Retrieved from <http://www.iucnredlist.org/> as accessed on 11 August 2010.
- Gelatt, T. S., Siniff, D. B., & Estes, J. A. (2002). Activity patterns and time budgets of the declining sea otter population at Amchitka Island, Alaska. *Journal of Wildlife Management*, 66, 29-39.
- Gentry, R. L. (2002). Northern Fur Seal. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 813-817). San Diego, CA: Academic Press.
- Gentry, R. L., Kooyman, G. L., & Goebel, M. E. (1986). Feeding and Diving Behavior of Northern Fur Seals. In Gentry, R. L. & Kooyman, G. L. (Eds.), *Fur Seals: Maternal Strategies on Land and at Sea* (pp. 61-78). Princeton: Princeton University Press.

- Gjertz, I., Lydersen, C., & Wiig, O. (2001). Distribution and Diving of Harbour Seals (*Phoca vitulina*) in Svalbard. *Polar Biology*, 24(3), 209-214.
- Goebel, M. E., Bengtson, J. L., Delong, R. L., Gentry, R. L., & Loughlin, T. R. (1991). Diving Patterns and Foraging Locations of Female Northern Fur Seals. *U S National Marine Fisheries Service Fishery Bulletin*, 89(2), 171-180.
- Goff, G. P., & Lien, J. (1988). Atlantic Leatherback Turtles, *Dermochelys coriacea*, in Cold Water off Newfoundland and Labrador. *Canadian Field-Naturalist*, 102(1), 1-5.
- Goldbogen, J. A., Calambokidis, J., Croll, D. A., Harvey, J. T., Newton, K. M., Oleson, E. M., Schorr, G., & Shadwick, R. E. (2008). Foraging Behavior of Humpback Whales: Kinematic and Respiratory Patterns Suggest a High Cost for a Lunge. *The Journal of Experimental Biology*, 211, 3712-3719.
- Goldbogen, J. A., Calambokidis, J., Croll, D. A., McKenna, M. F., Oleson, E., Potvin, J., Pyenson, N. D., Schorr, G., Shadwick, R. E., & Tershy, B. R. (2012). Scaling of lunge-feeding performance in rorqual whales: Mass-specific energy expenditure increases with body size and progressively limits diving capacity. *Functional Ecology*, 26, 216-226. doi: 10.1111/j.1365-2435.2011.01905.x.
- Goldbogen, J. A., Calambokidis, J., Friedlaender, A. S., Francis, J., DeRuiter, S. L., Stimpert, A. K., Falcone, E., & Southall, B. L. (2013). Underwater acrobatics by the world's largest predator: 360° rolling manoeuvres by lunge-feeding blue whales. *Biology Letters*, 9, 20120986. doi: 10.1098/rsbl.2012.0986.
- Goldbogen, J. A., Calambokidis, J., Shadwick, R. E., Oleson, E. M., McDonald, M. A., & Hildebrand, J. A. (2006). Kinematics of Foraging Dives and Lunge-Feeding in Fin Whales. *The Journal of Experimental Biology*, 209, 1231-1244.
- Green, G. A., Brueggeman, J. J., Grotefendt, R. A., Bowlby, C. E., Bonnell, M. L., & K.C. Balcomb III. (1992). *Cetacean Distribution and Abundance off Oregon and Washington, 1989-1990*. Los Angeles, California: Minerals Management Service. pp. 1-1 to 1-100.
- Guerrero, J. A. (1989). *Feeding Behavior of Gray Whales in Relation to Patch Dynamics of their Benthic Prey*. Master of Science, San José State University, San José, CA.
- Hain, J. H. W., Carter, G. R., Kraus, S. D., Mayo, C. A., & Winn, H. E. (1982). Feeding Behavior of the Humpback Whale, *Megaptera novaeangliae*, in the Western North Atlantic. *Fishery Bulletin*, 80(2), 259-268.
- Hakoyama, H., Le Boeuf, B. J., Naito, Y., & Sakamoto, W. (1994). Diving behavior in relation to ambient water temperature in northern elephant seals. *Canadian Journal of Zoology*, 72, 643-651.
- Hall, J. D. (1970). *Conditioning Pacific White Stripped Dolphins, Lagenorhynchus obliquidene, For Open Ocean Release*. (Naval Undersea Center Technical Publication 200). San Diego, CA: Naval Undersea Research and Development Center. p. 13.
- Hanser, S., Becker, E., Thorson, P., Zickel, M., & Hoopes, R. (2017). *U.S. Navy Marine Species Density Database Phase III for the Hawaii-Southern California Training and Testing Study Area*. Pearl Harbor, HI: U.S. Department of the Navy (DON), Commander U.S. Pacific Fleet (COMPACFLT), Naval Facilities Engineering Command Pacific (NAVFAC PAC), Naval Undersea Warfare Center (NUWC).
- Hastings, K. K., Frost, K. J., Simpkins, M. A., Pendleton, G. W., Swain, U. G., & Small, R. J. (2004). Regional Differences in Diving Behavior of Harbor Seals in the Gulf of Alaska. *Canadian Journal of Zoology*, 82, 1755-1773.

- Haug, T., Gjøsaeter, H., Lindstrøm, U., & Nilssen, K. T. (1995). Diet and Food Availability for North-East Atlantic Minke Whales (*Balaenoptera acutorostrata*), During the Summer of 1992. *ICES Journal of Marine Science*, 52(1), 77-86.
- Hays, G. C., Hobson, V. J., Metcalfe, J. D., Righton, D., & Sims, D. W. (2006). Flexible Foraging Movements of Leatherback Turtles Across the North Atlantic Ocean. *Ecology*, 87(10), 2647-2656.
- Hays, G. C., Houghton, J. D. R., Isaacs, C., King, R. S., Lloyd, C., & Lovell, P. (2004). First Records of Oceanic Dive Profiles for Leatherback Turtles, *Dermochelys coriacea*, Indicate Behavioural Plasticity Associated with Long-Distance Migration. *Animal Behaviour*, 67, 733-743.
- Hazen, E. L., Friedlaender, A. S., Thompson, M. A., Ware, C. R., Weinrich, M. T., Halpin, P. N., & Wiley, D. N. (2009). Fine-Scale Prey Aggregations and Foraging Ecology of Humpback Whales *Megaptera novaeangliae*. *Marine Ecology Progress Series*, 395, 75-89.
- Heaslip, S. G., Iverson, S. J., Bowen, W. D., & James, M. C. (2012). Jellyfish support high energy intake of leatherback sea turtles (*Dermochelys coriacea*): video evidence from animal-borne cameras. *PLoS ONE*, 7(3), e33259. doi: 10.1371/journal.pone.0033259.
- Heath, C. B. (2002). California, Galapagos, and Japanese Sea Lions. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 180-186). San Diego, CA: Academic Press.
- Heise, K. (1997). Diet and Feeding Behaviour of Pacific White-sided Dolphins (*Lagenorhynchus obliquidens*) as Revealed Through the Collection of Prey Fragments and Stomach Content Analyses. *Reports of the International Whaling Commission*, 47, 807-815.
- Herman, J. S., Kitchener, A. C., Baker, J. R., & Lockyer, C. (1994). The Most Northerly Record of Blainville's Beaked Whale, *Mesoplodon densirostris*, from the Eastern Atlantic. *Mammalia*, 58(4), 657-661.
- Heyning, J. E. (1989). Cuvier's Beaked Whale -- *Ziphius cavirostris* G. Cuvier, 1823. In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals: Volume 4: River Dolphins and the Larger Toothed Whales* (Vol. 4, pp. 289-308). New York: Academic Press.
- Heyning, J. E., & Perrin, W. F. (1994). Evidence for Two Species of Common Dolphins (Genus *Delphinus*) from the Eastern North Pacific. *Contributions in Science*, 442, 1-35.
- Hickmott, L. S. (2005). *Diving Behaviour and Foraging Behaviour and Foraging Ecology of Blainville's and Cuvier's Beaked Whales in the Northern Bahamas*. Master of Research in Environmental Biology (Master's), University of St. Andrews, Scotland.
- Hindell, M. A. (2002). Elephant Seals: *Mirounga angustirostris* and *M. leonina*. In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 370-373). San Diego, CA: Academic Press.
- Hobson, K. A., & Sease, J. L. (1998). Stable Isotope Analyses of Tooth Annuli Reveal Temporal Dietary Records: An Example Using Steller Sea Lions. *Marine Mammal Science*, 14(1), 116-129.
- Hobson, K. A., Sease, J. L., Merrick, R., L., & Piatt, J., F. (1997). Investigating Trophic Relationships of Pinnipeds in Alaska and Washington Using Stable Isotope Ratios of Nitrogen and Carbon. *Marine Mammal Science*, 13(1), 114-132.
- Horwood, J. (2002). Sei Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1069-1071). San Diego, CA: Academic Press.

- Houghton, J., Baird, R. W., Emmons, C. K., & Hanson, M. B. (2015). Changes in the occurrence and behavior of mammal-eating killer whales in southern British Columbia and Washington State, 1987–2010. *Northwest Science*.
- Houghton, J. D. R., Doyle, T. K., Davenport, J., Wilson, R. P., & Hays, G. C. (2008). The Role of Infrequent and Extraordinary Deep Dives in Leatherback Turtles (*Dermochelys coriacea*). *The Journal of Experimental Biology*, 211, 2566-2575.
- James, M. C., Davenport, J., & Hays, G. C. (2006a). Expanded Thermal Niche for a Diving Vertebrate: A Leatherback Turtle Diving into Near-Freezing Water. *Journal of Experimental Marine Biology and Ecology*, 335, 221-226.
- James, M. C., Eckert, S. A., & Myers, R. A. (2005a). Migratory and Reproductive Movements of Male Leatherback Turtles (*Dermochelys coriacea*). *Marine Biology*, 147, 845-853.
- James, M. C., Myers, R. A., & Ottensmeyer, C. A. (2005b). Behaviour of Leatherback Sea Turtles, *Dermochelys coriacea*, During the Migratory Cycle. *Proceedings of the Royal Society of London, Part B*, 272, 1547-1555.
- James, M. C., Ottensmeyer, C. A., Eckert, S. A., & Myers, R. A. (2006b). Changes in Diel Diving Patterns Accompany Shifts between Northern Foraging and Southward Migration in Leatherback Turtles. *Canadian Journal of Zoology*, 84, 754-765.
- Jefferson, T. A. (1988). *Phocoenoides dalli*. *Mammalian Species*, 319, 1-7.
- Jefferson, T. A. (2002). Dall's Porpoise: *Phocoenoides dalli*. In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 308-310). San Diego, CA: Academic Press.
- Jefferson, T. A., Fertl, D., Bolaños-Jiménez, J., & Zerbini, A. N. (2009). Distribution of Common Dolphins (*Delphinus spp.*) in the Western Atlantic Ocean: A Critical Re-examination. *Marine Biology*, 156, 1109-1124.
- Jefferson, T. A., & Newcomer, M. W. (1993a). *Lissodelphis borealis*. *Mammalian Species*, 425, 1-6.
- Jefferson, T. A., Newcomer, M. W., Leatherwood, S., & Van Waerebeek, K. (1994). Right Whale Dolphins—*Lissodelphis borealis* (Peale, 1848) and *Lissodelphis peronii* (Lacépède, 1804). In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals. Volume 5: The First Book of Dolphins* (Vol. 5, pp. 335-362). New York: Academic Press.
- Jefferson, T. A., & Newcomer, W. W. (1993b). *Lissodelphis borealis*. *Mammalian Species*, 425, 1-6.
- Jefferson, T. A., Smultea, M. A., Courbis, S. S., & Campbell, G. S. (2016). Harbor porpoise (*Phocoena phocoena*) recovery in the inland waters of Washington: Estimates of density and abundance from aerial surveys, 2013-2015. *Canadian Journal of Zoology*, 94. doi: 10.1139/cjz-2015-0236.
- Jefferson, T. A., Webber, M. A., & Pitman, R. L. (2008). *Marine Mammals of the World: A Comprehensive Guide to Their Identification*. San Diego, CA: Academic Press.
- Jensen, F. H., Perez, J. M., Johnson, M., Soto, N. A., & Madsen, P. T. (2011). Calling under pressure: short-finned pilot whales make social calls during deep foraging dives. *Proceedings of the Royal Society Biological Sciences*, 278, 3017-3025. doi: 10.1098/rspb.2010.2604.
- Johnson, M. P., & Aguilar de Soto, N. (2008a). Mesoplodon densirostris El Hierro, Canary Islands, Spain 10/11/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 11 October 2003 from <http://hdl.handle.net/1912/2367>.

- Johnson, M. P., & Aguilar de Soto, N. (2008b). Mesoplodon densirostris El Hierro, Canary Islands, Spain 10/25/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 25 October 2003 from <http://hdl.handle.net/1912/2366>.
- Johnson, M. P., & Sturlese, A. (2008a). Ziphius cavirostris Liguria, Italy 9/17/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 17 September 2003
- Johnson, M. P., & Sturlese, A. (2008b). Ziphius cavirostris Liguria, Italy 9/20/2003 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 20 September 2003
- Jones, M. L., & Swartz, S. L. (2002). Gray Whale (*Eschrichtius robustus*). In Perrin, W. F., Würsig, B. & Thewissen, H. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 524-536). San Diego, CA: Academic Press.
- Jonsen, I. D., Myers, R. A., & James, M. C. (2007). Identifying Leatherback Turtle Foraging Behavior from Satellite Telemetry using a Switching State-Space Model. *Ecology Progress Series*, 337, 255-264.
- Jorgensen, C., Lydersen, C., Brix, O., & Kovacs, K. M. (2001). Diving development in nursing harbour seal pups. *Journal of Experimental Biology*, 204, 3993-4004.
- Joyce, G. G., Sigurjonsson, J., & Vikingsson, G. (1990). Radio Tracking a Minke Whale (*Balaenoptera acutorostrata*) in Icelandic Waters for the Examination of Dive-time Patterns. *Reports of the International Whaling Commission*, 40, 357-361.
- Kajimura, H., & Loughlin, T. R. (1988). Marine Mammals in the Oceanic Food Web of the Eastern Subarctic Pacific. *Bulletin of the Ocean Research Institute, University of Tokyo*, 26, 187-223.
- Kasuya, T. (1986). Distribution and Behavior of Baird's Beaked Whales off the Pacific Coast of Japan. *The Scientific Reports of the Whales Research Institute*, 37, 61-83.
- Kasuya, T. (2002). Giant Beaked Whales. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 519-522). San Diego, CA: Academic Press.
- Kato, H. (2002). Bryde's Whales. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 171-177). San Diego, CA: Academic Press.
- Kawakami, T. (1980). A Review of Sperm Whale Food. *Scientific Reports of the Whales Research Institute*, 32, 199-218.
- Kenyon, K. W. (1981). Sea Otter - *Enhydra lutris*. In Ridgway, S. H. & Harrison, R. J. (Eds.), *Handbook of Marine Mammals* (Vol. 1: The Walrus, Sea Lions, Fur Seals and Sea Otter, pp. 209-224). San Diego, CA: Academic Press.
- Klatsky, L. J. (2004). *Movement and dive behavior of bottlenose dolphins (Tursiops truncatus) near the Bermuda Pedestal*. (masters), San Diego State University;Biology.
- Kooyman, G. L., Gentry, R. L., & Urquhart, D. L. (1976). Northern Fur Seal Diving Behavior: A New Approach to its Study. *Science*, 193(4251), 411-412.
- Kuhn, C. (2004). *Assessing the Feeding Behavior of California sea lions*. Santa Cruz, CA: University of California. p. 9.
- Kuhn, C. E. (2006). *Measuring At-Sea Feeding to Understand the Foraging Behavior of Pinnipeds*. PhD, University of California Santa Cruz.

- Kuhn, C. E., Johnson, D. S., Ream, R. R., & Gelatt, T. S. (2009). Advances in the Tracking of Marine Species: Using GPS Locations to Evaluate Satellite Track Data and a Continuous-Time Movement Model. *Marine Ecology Progress Series*, 393, 97-109.
- Kuhn, C. E., Tremblay, Y., Ream, R. R., & Gelatt, T. S. (2010). Coupling GPS tracking with dive behavior to examine the relationship between foraging strategy and fine-scale movements of northern fur seals. *Endangered Species Research*, 12(2), 125-139.
- Kvadsheim, P. H., Miller, P. J., Tyack, P. L., Sivle, L. D., Lam, F.-P. A., & Fahlman, A. (2012). Estimated tissue and blood N₂ levels and risk of decompression sickness in deep-, intermediate-, and shallow-diving toothed whales during exposure to naval sonar. *Frontiers in physiology*, 3.
- Laerm, J., Wenzel, F., Craddock, J. E., Weinand, D., McGurk, J., Harris, M. J., Early, G. A., Mead, J. G., Potter, C. W., & Barros, N. B. (1997). New Prey Species for Northwestern Atlantic Humpback Whales. *Marine Mammal Science*, 13(4), 705-711.
- Laidre, K. L., Heide-Jørgensen, M. P., Heagerty, P., Cossio, A., Bergström, B., & Simon, M. (2010). Spatial Associations between Large Baleen Whales and their Prey in West Greenland. *Marine Ecology Progress Series*, 402, 269-284.
- Laidre, K. L., & Jameson, R. J. (2006). Foraging Patterns and Prey Selection in an Increasing and Expanding Sea Otter Population. *Journal of Mammalogy*, 87(4), 799-807.
- Laidre, K. L., Jameson, R. J., Gurarie, E., Jeffries, S. J., & Allen, H. (2009). Spatial habitat use patterns of sea otters in coastal Washington. *Journal of Mammalogy*, 90, 906-917.
- Lander, M. E., Gulland, F. M. D., & DeLong, R. L. (2000). Satellite Tracking a Rehabilitated Guadalupe Fur Seal (*Arctocephalus townsendi*). *Aquatic Mammals*, 26(2), 137-142.
- Le Boeuf, B. J., Costa, D. P., Huntley, A. C., & Feldkamp, S. D. (1988). Continuous, Deep Diving in Female Northern Elephant Seals, *Mirounga angustirostris*. *Canadian Journal of Zoology*, 66(2), 446-458.
- Le Boeuf, B. J., Costa, D. P., Huntley, A. C., Kooyman, G. L., & Davis, R. W. (1986). Pattern and depth of dives in Northern elephant seals, *Mirounga angustirostris*. *The Zoological Society of London*, 208, 1-7.
- Le Boeuf, B. J., Crocker, D. E., Blackwell, S. B., Morris, P. A., & Thorson, P. H. (1993). Sex Differences in Diving and Foraging Behavior of Northern Elephant Seals. In: Boyd, I. L. (Ed.), *Marine Mammals: Advances in Behavioural and Population Biology* (Vol. 66, pp. 149-178). Oxford: Clarendon Press.
- Leatherwood, S., & Walker, W. A. (1979). The Northern Right Whale Dolphin *Lissodelphis borealis* Peale in the Eastern North Pacific. In: Winn, H. E. & Olla, B. L. (Eds.), *Behavior of Marine Animals: Current Perspectives in Research* (pp. 85-141). New York, NY: Plenum Press.
- LeDuc, R. G., Perrin, W. F., & Dizon, A. E. (1999). Phylogenetic Relationships Among the Delphinid Cetaceans Based on Full Cytochrome *b* Sequences. *Marine Mammal Science*, 15(3), 619-648.
- Linnenschmidt, M., Akamatsu, T., & Miller, A. (2013). Biosonar, dive, and foraging activity of satellite tracked harbor porpoises (*Phocoena phocoena*). *Marine Mammal Science*, 29(2), E77-E97. doi: 10.1111/j.1748-7692.2012.00592.x.
- Ljungblad, D. K., Moore, S. E., Clarke, J., & Bennett, J. C. (1987). *Distribution, Abundance, Behavior and Bioacoustics of Endangered Whales in the Alaskan Beaufort and Eastern Chukchi Seas, 1979-86*. (1177). San Diego, CA: Naval Ocean Systems Center (NOSC). p. 390.

- Loughlin, T. R. (2002). Steller's Sea Lion. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1181-1185). San Diego, CA: Academic Press.
- Loughlin, T. R., Perlov, A. S., Baker, J. D., Blokhin, S. A., & Makhnyr, A. G. (1998). Diving Behavior of Adult Female Steller Sea Lions in the Kuril Islands, Russia. *Biosphere Conservation*, 1(1), 21-31.
- Loughlin, T. R., Sterling, J. T., Merrick, R. L., Sease, J. L., & York, A. E. (2003). Diving Behavior of Immature Stellar Sea Lions (*Eumetopias jubatus*). *Fishery Bulletin*, 101(3), 566-582.
- Lowry, M. S., & Carretta, J. V. (1999). Market Squid (*Loligo opalescens*) in the Diet of California Sea Lions (*Zalophus californianus*) in Southern California (1981-1995). *CalCOFI Reports*, 40, 196-207.
- MacLeod, C. D., Santos, M. B., & Pierce, G. J. (2003a). Review of Data on Diets of Beaked Whales: Evidence of Niche Separation and Geographic Segregation. *Journal of the Marine Biological Association of the United Kingdom*, 83, 651-665.
- MacLeod, K., Simmonds, M. P., & Murray, E. (2003b). Summer Distribution and Relative Abundance of Cetacean Populations off North-West Scotland. *Journal of the Marine Biological Association of the United Kingdom*, 83, 1187-1192.
- Malcolm, C. D., & Duffus, D. A. (2000). Comparison of Subjective and Statistical Methods of Dive Classification Using Data from a Time-Depth Recorder Attached to a Gray Whale (*Eschrichtius robustus*). *Journal of Cetacean Research and Management*, 2(3), 177-182.
- Malcolm, C. D., Duffus, D. A., & Wischniowski, S. G. (1995). Small Scale Behaviour of Large Scale Subjects: Diving Behaviour of a Gray Whale (*Eschrichtius robustus*). *Western Geography*, 5/6, 35-44.
- Maravilla-Chavez, M. O., & Lowry, M. S. (1999). Incipient breeding colony of Guadalupe fur seals at isla Benito del este, Baja California, Mexico. *Marine Mammal Science*, 15(1), 239-241.
- Martin, A. R., & Clarke, M. R. (1986). The Diet of Sperm Whales (*Physeter macrocephalus*) Captured Between Iceland and Greenland. *Journal of the Marine Biological Association of the United Kingdom*, 66, 779-790.
- Mate, B. R., Palacios, D. M., Baker, C. S., Lagerquist, B. A., Irvine, L. M., Follett, T., Steel, D., Hayslip, C., & Winsori, M. H. (2016). *Baleen (Blue & Fin) Whale Tagging and Analysis in Support of Marine Mammal Monitoring Across Multiple Navy Training Areas. Preliminary Summary*. (Contract No. N62470-15-8006, Monitoring report). p. 60.
- McAlpine, D. F., Murison, L. D., & Hoberg, E. P. (1997). New Records for the Pygmy Sperm Whale, *Kogia breviceps* (Physeteridae) from Atlantic Canada with Notes on Diet and Parasites. *Marine Mammal Science*, 13(4), 701-704.
- Mead, J. G. (1989). Beaked Whales of the Genus -- *Mesoplodon*. In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 4: River Dolphins and the Larger Toothed Whales, pp. 349-430). London, United Kingdom: Academic Press.
- Mead, J. G., & Potter, C. W. (1990). Natural History of Bottlenose Dolphins Along the Central Atlantic Coast of the United States. In: Leatherwood, S. & Reeves, R. R. (Eds.), *The Bottlenose Dolphin* (pp. 165-195). New York: Academic Press, Inc.
- Melin, S. R., & DeLong, R. L. (1999). *At-Sea Distribution and Diving Behavior of California Sea Lion Females from San Miguel Island, California*. Seattle, WA: Alaska Fisheries Science Center, National Marine Fisheries Service, National Marine Mammal Laboratory. pp. 407-412.

- Melin, S. R., DeLong, R. L., & Siniff, D. B. (2008). The Effects of El Nino on the Foraging Behavior of Lactating California Sea Lions (*Zalophus californianus californianus*) During the Nonbreeding Season. *Canadian Journal of Zoology*, *86*, 192-206.
- Merrick, R. L. (1995). *The Relationship of the Foraging Ecology of Steller Sea Lions (Eumetopias jubatus) to their Population Decline in Alaska*. PhD, University of Washington, Seattle, WA.
- Merrick, R. L., & Loughlin, T. R. (1997). Foraging Behavior of Adult Female and Young-of-the-year Steller Sea Lions in Alaskan Waters. *Canadian Journal of Zoology*, *75*, 776-786.
- Merrick, R. L., Loughlin, T. R., Antonelis, G. A., & Hill, R. (1994). Use of Satellite-Linked Telemetry to Study Steller Sea Lion and Northern Fur Seal Foraging. *Polar Research*, *13*(1), 105-114.
- Miller, P., Antunes, R., Alves, A. C., Wensveen, P., Kvadsheim, P., Kleivane, L., Nordlund, N., Lam, F.-P., van IJsselmuide, S., Visser, F., Tyack, P. M., Patrick, Antunes, R., Alves, A. C., Wensveen, P., Kvadsheim, P., Kleivane, L., Nordlund, N., Lam, F.-P., van IJsselmuide, S., Visser, F., & Tyack, P. (2011). *The 3S Experiments: Studying the Behavioural Effects of Naval Sonar on Killer Whales (Orcinus orca), Sperm Whales (Physeter macrocephalus), and Long-finned Pilot Whales (Globicephala melas) in Norwegian Waters*.
- Miller, P. J. O. M., Shapiro, A. D., & Deecke, V. B. (2010). The diving behavior of mammal-eating killer whales (*Orcinus orca*): variations with ecological not physical factors. *Canadian Journal of Zoology*, *88*, 1103-1112.
- Minamikawa, S., Iwasaki, T., & Kishiro, T. (2007). Diving Behaviour of a Baird's Beaked Whale, *Berardius bairdii*, in the Slope Water Region of the western North Pacific: First Dive Records using a Data Logger. *Fisheries Oceanography*, *16*(6), 573-577.
- Minamikawa, S., Iwasaki, T., Tanaka, Y., Ryono, A., Noji, S., Sato, H., Kurosawa, S., & Kato, H. (2003, 17-21 March 2003). *Diurnal Pattern of Diving Behavior in Striped Dolphins, Stenella coeruleoalba*. Paper presented at the International Symposium on Bio-logging Science, Tokyo, Japan.
- Mintzer, V. J., Gannon, D. P., Barros, N. B., & Read, A. J. (2008). Stomach contents of mass-stranded short-finned pilot whales (*Globicephala macrorhynchus*). *Marine Mammal Science*, *24*(2), 290-302.
- Miyazaki, N., Kuramochi, T., & Amano, M. (1991). Pacific White-sided Dolphins (*Lagenorhynchus obliquidens*) Off Northern Hokkaido. *Memoirs of the National Science Museum, Tokyo*, *24*, 131-139.
- Mizroch, S. A., Rice, D. W., & Breiwick, J. M. (1984). The Sei Whale, *Balaenoptera borealis*. *Marine Fisheries Review*, *46*(4), 25-29.
- Morton, A. (2000). Occurrence, Photo-Identification and Prey of Pacific White-Sided Dolphins (*Lagenorhynchus obliquidens*) in the Broughton Archipelago, Canada 1984-1998. *Marine Mammal Science*, *16*(1), 80-93.
- Myers, A. E., & Hays, G. C. (2006). Do Leatherback Turtles *Dermochelys coriacea* Forage During the Breeding Season? A Combination of Data-Logging Devices Provide New Insights. *Marine Ecology Progress Series*, *322*, 259-267.
- Naito, Y., Le Boeuf, B. J., Asaga, T., & Huntley, A. C. (1989). Long Term Diving Records of an Adult Female Northern Elephant Seal. *Antarctic Record*, *33*(1), 1-9.
- National Marine Fisheries Service. (2015, 16 January 2015). Bottlenose Dolphin (*Tursiops truncatus*). *Protected Resources-Species* Retrieved Retrieved 16 January 2015 from

- <http://www.nmfs.noaa.gov/pr/species/mammals/dolphins/bottlenose-dolphin.html> as accessed on 2 September 2016.
- National Marine Fisheries Service (NMFS), & United States Fish and Wildlife Service (USFWS). (1993). *Recovery Plan for Hawksbill Turtles in the U.S. Caribbean Sea, Atlantic Ocean, and Gulf of Mexico*. St Petersburg, FL: National Marine Fisheries Service (NMFS). p. 55.
- Nerini, M. (1984). A Review of Gray Whale Feeding Ecology. In: Jones, M. L., Swartz, S. L. & Leatherwood, S. (Eds.), *The Gray Whale (Eschrichtius robustus)* (pp. 423-448). Orlando, FL: Academic Press.
- Network, O. (2017, 2017/09//). Whale sightings for Puget Sound downloaded from the Orca Network webpages for 2003-2017 Retrieved Retrieved 2017/09//from http://www.orcanetwork.org/Archives/index.php?categories_file=Sightings%20Archives%20Home as accessed on September 2017.
- Neumann, D. R., & Orams, M. B. (2003). Feeding Behaviours of Short-beaked Common Dolphins, *Delphinus delphis*, in New Zealand. *Aquatic Mammals*, 29(1), 137-149.
- Newby, T. C. (1975). A Sea Otter (*Enhydra lutris*) Food Dive Record. *The Murrelet*, 56(1), 7.
- Noren, S. R., & Williams, T. M. (2000). Body Size and Skeletal Muscle Myoglobin of Cetaceans: Adaptations for Maximizing Dive Duration. *Comparative Biochemistry and Physiology Part A*, 126, 181-191.
- Nøttestad, L., Fernö, A., & Axelsen, B. E. (2002). Digging in the Deep: Killer Whales' Advanced Hunting Tactic. *Polar Biology*, 25, 939-941.
- Nøttestad, L., Fernö, A., Mackinson, S., Pitcher, T., & Misund, O. A. (2002). How Whales Influence Herring School Dynamics in a Cold-Front Area of the Norwegian Sea. *ICES Journal of Marine Science*, 59(2), 393-400.
- Ohizumi, H., Kuramochi, T., Amano, M., & Miyazaki, N. (2000). Prey Switching of Dall's Porpoise *Phocoenoides dalli* with Population Decline of Japanese Pilchard *Sardinops melanostictus* Around Hokkaido, Japan. *Marine Ecology Progress Series*, 200, 265-275.
- Ohizumi, H., Yoshioka, M., Mori, K., & Miyazaki, N. (1998). Stomach Contents of Common Dolphins (*Delphinus delphis*) in the Pelagic Western North Pacific. *Marine Mammal Science*, 14(4), 835-844.
- Øien, N. (1990). Sightings Surveys in the Northeast Atlantic in July 1988: Distribution and Abundance of Cetaceans. *Reports of the International Whaling Commission*, 40, 499-511.
- Olesiuk, P. F. (1993). Annual Prey Consumption by Harbor Seals (*Phoca vitulina*) in the Strait of Georgia, British Columbia. *Fishery Bulletin*, 91(3), 491-515.
- Oleson, E. M., Calambokidis, J., Burgess, W. C., McDonald, M. A., LeDuc, C. A., & Hildebrand, J. A. (2007). Behavioral Context of Call Production by Eastern North Pacific Blue Whales. *Marine Ecology Progress Series*, 330, 269-284.
- Omura, H. (1962). Further Information on Bryde's Whale from the Coast of Japan. *Scientific Reports of the Whales Research Institute*, 16, 7-18.
- Otani, S., Naito, Y., Kato, A., & Kawamura, A. (2000). Diving Behavior and Swimming Speed of a Free-Ranging Harbor Porpoise, *Phocoena phocoena*. *Marine Mammal Science*, 16(4), 811-814.

- Otani, S., Naito, Y., Kawamura, A., Kawasaki, M., Nishiwaki, S., & Kato, A. (1998). Diving Behavior and Performance of Harbor Porpoises, *Phocoena phocoena*, in Funka Bay, Hokkaido, Japan. *Marine Mammal Science*, 14(2), 209-220.
- Panigada, S., Zanardelli, M., Canese, S., & Jahoda, M. (1999, 28 November - 03 December 1999). *Deep Diving Performances of Mediterranean Fin Whales*. Paper presented at the 13th Biennial Conference on the Biology of Marine Mammals, Wailea, HI.
- Payne, P. M., & Selzer, L. A. (1989). The Distribution, Abundance, and Selected Prey of the Harbor Seal, *Phoca vitulina concolor*, in Southern New England. *Marine Mammal Science*, 5(2), 173-192.
- Perrin, W. F., & Hohn, A. A. (1994). Pantropical Spotted Dolphin--*Stenella attenuata*. In: Ridgway, S. H. & Harrison, S. R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 71-98). New York, NY: Academic Press.
- Perrin, W. F., Wilson, C. E., & Archer II, F. I. (1994). Striped Dolphin--*Stenella coeruleoalba* (Meyen, 1833). In: Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 5: The First Book of Dolphins, pp. 129-159). New York, NY: Academic Press.
- Pitman, R. L. (2002). Mesoplodont Whales. In: Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 738-742). San Diego, CA: Academic Press.
- Pitman, R. L., & Ensor, P. (2003). Three Forms of Killer Whales (*Orcinus orca*) in Antarctic Waters. *Journal of Cetacean Research and Management*, 5(2), 131-139.
- Plön, S. (2004). *The Status and Natural History of Pygmy (Kogia breviceps) and Dwarf (K. sima) Sperm Whales off Southern Africa*. PhD in Zoology (Ph.D Dissertation), Rhodes University, Grahamstown, South Africa.
- Ponganis, P. J., Gentry, R. L., Ponganis, E. P., & Ponganis, K. V. (1992). Analysis of Swim Velocities During Deep and Shallow Dives of Two Northern Fur Seals *Callorhinus ursinus*. *Marine Mammal Science*, 8(1), 69-75.
- Porras-Peters, H., Auriolles-Gamboa, D., Cruz-Escalona, V. H., & Koch, P. L. (2008). Trophic Level and Overlap of Sea Lions (*Zalophus californianus*) in the Gulf of California, Mexico. *Marine Mammal Science*, 24(3), 554-576.
- Pusineri, C., Magnin, V., Meynier, L., Spitz, J., Hassani, S., & Ridoux, V. (2007). Food and Feeding Ecology of the Common Dolphin (*Delphinus delphis*) in the Oceanic Northeast Atlantic and Comparison with its Diet in Neritic Areas. *Marine Mammal Science*, 23(1), 30-47.
- Rankin, S., Oswald, J., Barlow, J., & Lammers, M. (2007). Patterned Burst-Pulse Vocalizations of the Northern Right Whale Dolphin, *Lissodelphis borealis*. *Journal of the Acoustical Society of America*, 121(12), 1213-1218.
- Ream, R. R., Sterling, J. T., & Loughlin, T. R. (2005). Oceanographic Features Related to Northern Fur Seal Migratory Movements. *Deep Sea Research II*, 52, 823-843.
- Recchia, C. A., & Read, A. J. (1989). Stomach Contents of Harbour Porpoises, *Phocoena phocoena* (L.), from the Bay of Fundy. *Canadian Journal of Zoology*, 67, 2140-2146.
- Reeves, R. R., Stewart, B. S., Clapham, P. J., & Powell, J. A. (2002). *Guide to Marine Mammals of the World*. New York, NY: Chanticleer Press Inc.

- Rehberg, M. J., Andrews, R. D., Swain, U. G., & Calkins, D. G. (2009). Foraging Behavior of Adult Female Steller Sea Lions During the Breeding Season in Southeast Alaska. *Marine Mammal Science*, 25(3), 588-604.
- Riedman, M. (1990). *The Pinnipeds: Seals, Sea Lions, and Walruses*. Berkley, CA: University of California Press.
- Riedman, M. L., & Estes, J. A. (1990). *The Sea Otter (Enhydra lutris): Behavior, Ecology, and Natural History*. (Biological Report 90(14)). Washington, DC: United States Department of the Interior. p. 33.
- Ringelstein, J., Pusineri, C., Hassani, S., Meynier, L., Nicolas, R., & Ridoux, V. (2006). Food and Feeding Ecology of the Striped Dolphin, *Stenella coeruleoalba*, in the Oceanic Waters of the North-east Atlantic. *Journal of the Marine Biological Association of the United Kingdom*, 86, 909-918.
- Robertson, K. M., & Chivers, S. J. (1997). Prey Occurrence in Pantropical Spotted Dolphins, *Stenella attenuata*, from the Eastern Tropical Pacific. *Fishery Bulletin*, 95, 334-348.
- Robinson, P. W., Costa, D. P., Crocker, D. E., Gallo-Reynoso, J. P., Champagne, C. D., Fowler, M. A., Goetsch, C., Goetz, K. T., Hassrick, J. L., Huckstadt, L. A., Kuhn, C. E., Maresh, J. L., Maxwell, S. M., McDonald, B. I., Peterson, S. H., Simmons, S. E., Teutschel, N. M., Villegas-Amtmann, S., & Yoda, K. (2012). Foraging Behavior and Success of a Mesopelagic Predator in the Northeast Pacific Ocean: Insights from a Data-Rich Species, the Northern Elephant Seal. *PloS one*, 7(5). doi: 10.1371/journal.pone.0036728.
- Rosel, P. E., Dizon, A. E., & Heyning, J. E. (1994). Genetic Analysis of Sympatric Morphotypes of Common Dolphins (genus *Delphinus*). *Marine Biology*(119), 159-167.
- Ross, G. J. B. (1979). Records of Pygmy and Dwarf Sperm Whales, genus *Kogia*, from Southern Africa, with Biological Notes and Some Comparisons. *Annals of the Cape Provincial Museums (Natural History)*, 11(14), 259-327.
- Roszbach, K. A., & Herzing, D. L. (1997). Underwater Observations of Benthic-Feeding Bottlenose Dolphins (*Tursiops truncatus*) Near Grand Bahama Island, Bahamas. *Marine Mammal Science*, 13(3), 498-504.
- Ruchonnet, D., Boutoute, M., Guinet, C., & Mayzaud, P. (2006). Fatty Acid Composition of Mediterranean Fin Whale *Balaenoptera physalus* Blubber with Respect to Body Heterogeneity and Trophic Interaction. *Marine Ecology Progress Series*, 311, 165-174.
- Sale, A., Luschi, P., Mencacci, R., Lambardi, P., Hughes, G. R., Hays, G. C., Benvenuti, S., & Papi, F. (2006). Long-Term Monitoring of Leatherback Turtle Diving Behaviour During Oceanic Movements. *Journal of Experimental Marine Biology and Ecology*, 328, 197-210.
- Santos, M. B., Pierce, G. J., Herman, J., Lopez, A., Guerra, A., Mente, E., & Clarke, M. R. (2001). Feeding Ecology of Cuvier's Beaked Whale (*Ziphius cavirostris*): A Review with New Information on the Diet of this Species. *Journal of the Marine Biological Association of the United Kingdom*, 81, 687-694.
- Santos, M. B., Pierce, G. J., Lopez, A., Reid, R. J., Ridoux, V., & Mente, E. (2006). Pygmy Sperm Whales *Kogia breviceps* in the Northeast Atlantic: New Information on Stomach Contents and Strandings. *Marine Mammal Science*, 22(3), 600-616.

- Sasaki, T., Nikaido, M., Hamilton, H., Goto, M., Kato, H., Kanda, N., Pastene, L. A., Cao, Y., Fordyce, R. E., Hasbgawa, M., & Okada, N. (2005). Mitochondrial Phylogenetics and Evolution of Mysticete Whales. *Systematic Biology*, 54(1), 77-90.
- Sasaki, T., Nikaido, M., Wada, S., Yamada, T. K., Cao, Y., Hasegawa, M., & Okada, N. (2006). *Balaenoptera omurai* is a Newly Discovered Baleen Whale that Represents an Ancient Evolutionary Lineage. *Molecular Phylogenetics and Evolution*, 41, 40-52.
- Saulitis, E., Matkin, C., Barrett-Lennard, L., Heise, K., & Ellis, G. (2000). Foraging Strategies of Sympatric Killer Whale (*Orcinus orca*) Populations in Prince William Sound, Alaska. *Marine Mammal Science*, 16(1), 94-109.
- Schoenherr, J. R. (1991). Blue whales Feeding on High Concentrations of Euphausiids Around Monterey Submarine Canyon. *Canadian Journal of Zoology*, 69, 583-594.
- Schorr, G. S., Falcone, E. A., Moretti, D. J., & Andrews, R. D. (2014). First Long-Term Behavioral Records from Cuvier's Beaked Whales (*Ziphius cavirostris*) Reveal Record-Breaking Dives. *PLOS ONE*, 9(3), e92633.
- Schusterman, R. J. (1981). Steller Sea Lion *Eumetopias jubatus* (Schreber, 1776). In: Ridgway, S. & Harrison, R. (Eds.), *Handbook of Marine Mammals: Vol 1. Walrus, Sea Lions, Fur Seals and Sea Otters* (pp. 119-142). New York, NY: Academic Press.
- Scott, M. D., & Chivers, S. J. (2009). Movements and Diving Behavior of Pelagic Spotted Dolphins. *Marine Mammal Science*, 25(1), 137-160.
- Scott, M. D., Hohn, A. A., Westgate, A. J., Nicolas, J. R., Whitaker, B. R., & Campbell, W. J. (2001). A note on the release and tracking of a rehabilitated pygmy sperm whale (*Kogia breviceps*). *Journal of Cetacean Research and Management*, 3(1), 87-94.
- Sears, R. (2002). Blue Whale. In: Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 112-116). San Diego, CA: Academic Press.
- Sears, R., & Perrin, W. F. (2008). Blue Whale, 2nd. In *Encyclopedia of Marine Mammals* (pp. 120-124). San Diego, CA: Academic Press.
- Selzer, L. A., & Payne, P. M. (1988). The Distribution of White-Sided (*Lagenorhynchus acutus*) and Common Dolphins (*Delphinus delphis*) vs. Environmental Features of the Continental Shelf of the Northeastern United States. *Marine Mammal Science*, 4(2), 141-153.
- Shane, S. H. (1990). Behavior and Ecology of the Bottlenose Dolphin at Sanibel Island, Florida. In: Leatherwood, S. & Reeves, R. R. (Eds.), *The Bottlenose Dolphin* (pp. 245-265). New York, NY: Academic Press, Inc.
- Shapiro, A. D. (2008). *Orchestration: The Movement and Vocal Behavior of Free-Ranging Norwegian Killer Whales (Orcinus orca)*. Doctor of Philosophy (Doctrine), Massachusetts Institute of Technology and the Woods Hole Oceanographic Institution, Woods Hole, MA.
- Siciliano, S., Santos, M. C. d. O., Vicente, A. F. C., Alvarenga, F. S., Zampirolli, E., Brito Jr., J. L., Azevedo, A. F., & Pizzorno, J. L. A. (2004). Strandings and Feeding Records of Bryde's Whales (*Balaenoptera edeni*) in South-Eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom*, 84, 857-859.
- Sinclair, E. H. (1994). Prey of Juvenile Northern Elephant Seals (*Mirounga angustirostris*) in the Southern California Bight. *Marine Mammal Science*, 10(2), 230-239.

- Sivle, L. D., Kvadsheim, P. H., Fahlman, A., Lam, F. P. A., Tyack, P. L., & Miller, P. J. O. (2012). Changes in dive behavior during naval sonar exposure in killer whales, long-finned pilot whales, and sperm whales. *Frontiers in Physiology*, 3(Article 400), 1-11. doi: 10.3389/fphys.2012.00400.
- Skinner, J. P., Burkanov, V. N., & Andrews, R. D. (2012). Influence of environment, morphology, and instrument size on lactating northern fur seal *Callorhinus ursinus* foraging behavior on the Lovushki Islands, Russia. *Marine Ecology Progress Series*, 471, 293-308.
- Smith, R. J., & Read, A. J. (1992). Consumption of Euphausiids by Harbour Porpoise (*Phocoena phocoena*) Calves in the Bay of Fundy. *Canadian Journal of Zoology*, 70, 1629-1632.
- Smultea, M. A., Lomac-MacNair, K., Campbell, G., Courbis, S. S., & Jefferson, T. A. (2017). *Aerial surveys of marine mammals conducted in the inland Puget Sound waters of Washington, Summer 2013–Winter 2016 Final Report*. Preston, WA: Department of the Navy, Commander U.S. Pacific Fleet, Naval Sea Systems Command, Smultea Environmental Sciences.
- Soto, N. A., Johnson, M. P., Madsen, P. T., Díaz, F., Domínguez, I., Brito, A., & Tyack, P. (2008). Cheetahs of the Deep Sea: Deep Foraging Sprints in Short-Finned Pilot Whales off Tenerife (Canary Islands). *Journal of Animal Ecology*, 77(5), 936-947.
- Southall, B., Calambokidis, J., Barlow, J., Moretti, D., Friedlaender, A., Stimpert, A., Douglas, A., Southall, B., Arranz, P., DeRuiter, S., Hazen, E., Goldbogen, J., Falcone, E., & Schorr, G. (2014). *Biological and behavioral response studies of marine mammals in southern California, 2013 ("SOCAL-13"), final project report*. Southall Environmental Associates, Inc. p. 54.
- Stelle, L. L., Megill, W. M., & Kinzel, M. R. (2008). Activity Budget and Diving Behavior of Gray Whales (*Eschrichtius robustus*) in Feeding Grounds Off Coastal British Columbia. *Marine Mammal Science*, 24(3), 462-478.
- Sterling, J. T., & Ream, R. R. (2004). At-Sea Behavior of Juvenile Male Northern Fur Seals (*Callorhinus ursinus*). *Canadian Journal of Zoology*, 82(10), 1621-1637.
- Stewart, B. S., Harvey, J., & Yochem, P. K. (2001). Post-Release Monitoring and Tracking of a Rehabilitated California Gray Whale. *Aquatic Mammals*, 27(3), 294-300.
- Stewart, B. S., & Huber, H. R. (1993). *Mirounga angustirostris*. *Mammalian Species*(1-10).
- Stewart, B. S., Yochem, P. K., DeLong, R. L., & Antonelis, G. A. (1987). *Interactions between Guadalupe fur seals and California sea lions at San Nicolas and San Miguel Islands, California*. Paper presented at the Status, Biology and Ecology of Fur Seals, Cambridge, England.
- Stimpert, A. K., DeRuiter, S., Southall, B., Moretti, D., Falcone, E., Goldbogen, J., Friedlaender, A., Schorr, G., & Calambokidis, J. (2014). Acoustic and foraging behavior of a Baird's beaked whale, *Berardius bairdii*, exposed to simulated sonar. *Scientific Reports*, 4. doi: 10.1038/srep07031.
- Stockin, K. A., Fairbairns, R. S., Parsons, E. C. M., & Sims, D. W. (2001). Effects of Diel and Seasonal Cycles on the Dive Duration of the Minke Whale (*Balaenoptera acutorostrata*). *Journal of the Marine Biological Association of the United Kingdom*, 81(1), 189-190.
- Straley, J. M., Moran, J. R., Boswell, K. M., Vollenweider, J. J., Heintz, R. A., Quinn li, T. J., Witteveen, B. H., & Rice, S. D. (2018). Seasonal presence and potential influence of humpback whales on wintering Pacific herring populations in the Gulf of Alaska. *Deep Sea Research Part II: Topical Studies in Oceanography*, 147. doi: 10.1016/j.dsr2.2017.08.008.

- Stroud, R. K., Fiscus, C. H., & Kajimura, H. (1981). Food of the Pacific White-Sided Dolphin, *Lagenorhynchus obliquidens*, Dall's Porpoise, *Phocoenoides dalli*, and Northern Fur Seal, *Callorhinus ursinus*, off California and Washington. *Fishery Bulletin*, 78(4), 951-959.
- Swartz, S. L. (1986). Gray Whale Migratory, Social and Breeding Behavior. In: Donovan, G. P. (Ed.), *Behaviour of Whales in Relation to Management: Incorporating the Proceedings of a Workshop held in Seattle, WA, 19-23 April 1982* (Vol. Special Issue 8, pp. 207-229). Cambridge: International Whaling Commission.
- Teilmann, J., Christiansen, C. T., Kjellerup, S., Dietz, R., & Nachman, G. (2013). Geographic, seasonal, and diurnal surface behavior of harbor porpoises *Marine Mammal Science*, 29(2), E60-E76. doi: 10.1111/j.1748-7692.2012.00597.x.
- Teloni, V., Mark, J. P., Patrick, M. J. O., & Peter, M. T. (2008). Shallow Food for Deep Divers: Dynamic Foraging Behavior of Male Sperm Whales in a High Latitude Habitat. *Journal of Experimental Marine Biology and Ecology*, 354, 119-131.
- The Whale Museum. (2017a). *Orca Sightings Access Database 2006-2016 provided by Jennifer Olson on 9/8/2017 from The Whale Museum*. Friday Harbor, WA: The Whale Museum.
- The Whale Museum. (2017b). *Orca Sightings Access Database 2006-2016 provided by Jennifer Olson on 9/13/2017 from The Whale Museum*. Friday Harbor, WA: The Whale Museum.
- Thomas, K., Harvey, J. T., Goldstein, T., Barakos, J., & Gulland, F. (2010). Movement, dive behavior, and survival of California sea lions (*Zalophus californianus*) posttreatment for domoic acid toxicosis. *Marine Mammal Science*, 26(1), 36-52.
- Thometz, N. M., Staedler, M. M., Tomoleoni, J. A., Bodkin, J. L., Bentall, G. B., & Tinker, M. T. (2016). Trade-offs between energy maximization and parental care in a central place forager, the sea otter. *Behavioral Ecology*, 27(5), 1552-1566.
- Threlfall, W. (1978). First Record of the Atlantic Leatherback Turtle (*Dermochelys coriacea*) from Labrador. *Canadian Field-Naturalist*, 92(3), 287.
- Tollit, D. J., Heaslip, S. G., & Trites, A. W. (2004). Sizes of Walleye Pollock (*Theragra chalcogramma*) Consumed by the Eastern Stock of Steller Sea Lions (*Eumetopias jubatus*) in Southeast Alaska from 1994 to 1999. *Fishery Bulletin*, 102(3), 522-532.
- Tyack, P. (2010). Mesoplodon denirostris Andros Island, Bahamas 10/23/2006 Animal a Depth Profile. In The DTAG Project, W. H. O. I. (Ed.). Retrieved Retrieved 23 October 2006 from <http://hdl.handle.net/1912/3650>.
- Tyack, P. L., Johnson, M., Soto, N. A., Sturlese, A., & Madsen, P. T. (2006). Extreme Diving of Beaked Whales. *The Journal of Experimental Biology*, 209, 4238-4253.
- U.S. Department of the Navy. (2018). *Quantifying Acoustic Impacts on Marine Mammals and Sea Turtles: Methods and Analytical Approach for Phase III Training and Testing*. Newport, RI: Naval Undersea Warfare Center, Division Newport, Naval Warfare Systems Command Marine Mammal Program.
- Vikingsson, G. A. (1997). Feeding of Fin Whales (*Balaenoptera physalus*) off Iceland - Diurnal and Seasonal Variation and Possible Rates. *Journal of Northwest Atlantic Fishery Science*, 22, 77-89.
- Waerebeek, K. V., & Würsig, B. (2002). Pacific White-Sided Dolphin and Dusky Dolphin. In: Perrin, W. F., Würsig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 859-861). San Diego, CA: Academic Press.

- Walker, W. A. (1996). Summer Feeding Habits of Dall's Porpoise, *Phocoenoides dalli*, in the Southern Sea of Okhotsk. *Marine Mammal Science*, 12(2), 167-182.
- Walker, W. A., Hanson, M. B., Baird, R. W., & Guenther, T. J. (1998). Food Habits of the Harbor Porpoise, *Phocoena phocoena*, and Dall's Porpoise, *Phocoenoides dalli*, in the Inland Waters of British Columbia and Washington. In Alaska Fisheries Science Center, A. (Ed.), *Marine Mammal Protection Act and Endangered Species Act Implementation Program 1997, AFSC Processed Report 98-10* (pp. 14): National Marine Fisheries Service.
- Walker, W. A., & Jones, L. L. (1994). Food Habits of Northern Right Whale Dolphin, Pacific White-Sided Dolphin, and Northern Fur Seal Caught in the High Seas Driftnet Fisheries of the North Pacific Ocean, 1990. *International North Pacific Fisheries Bulletin*, 53(2), 285-295.
- Walker, W. A., Mead, J. G., & Brownell Jr., R. L. (2002). Diets of Baird's Beaked Whales, *Berardius bairdii*, in the Southern Sea of Okhotsk and off the Pacific Coast of Honshu, Japan. *Marine Mammal Science*, 18(4), 902-919.
- Wallace, B. P., Tiwari, M., & Girondot, M. (2013). *Dermochelys coriacea*. . *The IUCN Red List of Threatened Species*. Retrieved from www.iucnredlist.org
- Wang, M.-C., Walker, W. A., Shao, K.-T., & Chou, L.-S. (2002). Comparative Analysis of the Diets of Pygmy Sperm Whales and Dwarf Sperm Whales in Taiwanese Waters. *Acta Zoologica Taiwanica*, 13(2), 53-62.
- Wang, M.-C., Walker, W. A., Shao, K.-T., & Chou, L.-S. (2003). Feeding Habits of the Pantropical Spotted Dolphin, *Stenella attenuata*, off the Eastern Coast of Taiwan. *Zoological Studies*, 42(2), 368-378.
- Waring, G. T., Josephson, E., Fairfield, C. P., & Maze-Foley, K. (2006). *U.S. Atlantic and Gulf of Mexico Marine Mammal Stock Assessments 2005*. Woods Hole, MA: U.S. Department of Commerce National Oceanic and Atmospheric Administration National Marine Fisheries Service. p. 358.
- Watwood, S. L., Miller, P. J. O., Johnson, M., Madsen, P. T., & Tyack, P. L. (2006). Deep-Diving Foraging Behaviour of Sperm Whales (*Physeter macrocephalus*). *Journal of Animal Ecology*, 75, 814-825.
- Weir, C. R. (2007). Occurrence and Distribution of Cetaceans off Northern Angola, 2004/05. *Journal of Cetacean Research and Management*, 9(3), 225-239.
- Weise, M. J. (2006). *Foraging Ecology of Male California Sea Lion (Zalophus californianus): Movement, Diving and Foraging Behavior, and Diving Capacity*. Doctor of Philosophy (Doctrine), University of California, Santa Cruz, CA.
- Wells, R. S., Fougères, E. M., Cooper, A. G., Stevens, R. O., Brodsky, M., Lingenfelter, R. G., Dold, C., & Douglas, D. C. (2013). Movements and dive patterns of short-finned pilot whales (*Globicephala macrorhynchus*) released from a mass stranding in the Florida Keys. *Aquatic Mammals*, 39(1), 61-72.
- Wells, R. S., Manire, C. A., Byrd, L., Smith, D. R., Gannon, J. G., Fauquier, D., & Mullin, K. D. (2009). Movements and Dive Patterns of a Rehabilitated Risso's Dolphin, *Grampus griseus*, in the Gulf of Mexico and the Atlantic Ocean. *Marine Mammal Science*, 25(2), 420-429.
- Wells, R. S., & Scott, M. D. (1999). Bottlenose Dolphin - *Tursiops truncatus* (Montagu, 1821). In Ridgway, S. H. & Harrison, R. (Eds.), *Handbook of Marine Mammals* (Vol. 6: The Second Book of Dolphins and the Porpoises, pp. 137-182). San Diego, CA: Academic Press.

- Wells, R. S., & Scott, M. D. (2002). Bottlenose Dolphins, *Tursiops truncatus* and *T. aduncus*. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 122-128). San Diego, CA: Academic Press.
- Westgate, A. J., & Read, A. J. (1998). Applications of New Technology to the Conservation of Porpoises. *Marine Technology Science Journal*, 32(1), 70-81.
- Westgate, A. J., Read, A. J., Berggren, P., Koopman, H. N., & Gaskin, D. E. (1995). Diving Behaviour of Harbour Porpoises, *Phocoena phocoena*. *Canadian Journal of Fisheries and Aquatic Sciences*, 52, 1064-1073.
- Whitehead, H. (2002). Sperm Whale. In Perrin, W. F., Wursig, B. & Thewissen, J. G. M. (Eds.), *Encyclopedia of Marine Mammals* (pp. 1165-1172). San Diego, CA: Academic Press.
- Wolt, R. C. (2014). *Activity budget, field metabolic rate, and foraging ecology of female sea otters (Enhydra lutris kenyoni) with dependent pups in Alaska*. Doctor.
- Wolt, R. C., Gelwick, F. P., Weltz, F., & Davis, R. W. (2012). Foraging behavior and prey of sea otters in a soft- and mixed-sediment benthos in Alaska. *Mammalian Biology*, 77, 271-280.
- Womble, J. N., Blundell, G. M., Gende, S. M., Horning, M., Sigler, M. F., & Csepp, D. J. (2014). Linking marine predator diving behavior to local prey fields in contrasting habitats in a subarctic glacial fjord. *Marine Biology*, 161, 1361-1374.
- Woodward, B. (2006). *Locomotory Strategies, Dive Dynamics, and Functional Morphology of the Mysticetes: Using Morphometrics, Osteology, and DTAG Data to Compare Swim Performance in Four Species*. (Doctor of Philosophy), The University of Maine, Orono, ME.
- Woodward, B. L., & Winn, J. P. (2006). Apparent Lateralized Behavior in Gray Whales Feeding off the Central British Columbia Coast. *Marine Mammal Science*, 22(1), 64-73.
- Wursig, B., Wells, R. S., & Croll, D. A. (1986). Behavior of Gray Whales Summering near St. Lawrence Island, Bering Sea. *Canadian Journal of Zoology*, 64, 611-621.
- Yeates, L. C., Williams, T. M., & Fink, T. L. (2007). Diving and Foraging Energetics of the Smallest Marine Mammal, the Sea Otter (*Enhydra lutris*). *Journal of Experimental Biology*, 210, 1960-1970.

This page left intentionally blank.

INITIAL DISTRIBUTION LIST

Addressee

No. of Copies

Defense Technical Information Center

1